

Coming to Grips With Complex Disorders: Genetic Risk Prediction in Bipolar Disorder Using Panels of Genes Identified Through Convergent Functional Genomics

S.D. Patel,¹ H. Le-Niculescu,¹ D.L. Koller,² S.D. Green,¹ D.K. Lahiri,¹ F.J. McMahon,³ J.I. Nurnberger Jr,¹ and A.B. Niculescu III^{1,4*}

¹Department of Psychiatry, Indiana University School of Medicine, Indianapolis, Indiana

²Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, Indiana

³Mood and Anxiety Disorders Branch, NIMH, Bethesda, Maryland

⁴Indianapolis VA Medical Center, Indianapolis, Indiana

Received 26 January 2010; Accepted 19 February 2010

We previously proposed and provided proof of principle for the use of a complementary approach, convergent functional genomics (CFG), combining gene expression and genetic data, from human and animal model studies, as a way of mining the existing GWAS datasets for signals that are there already, but did not reach significance using a genetics-only approach [Le-Niculescu et al., 2009b]. CFG provides a fit-to-disease prioritization of genes that leads to generalizability in independent cohorts, and counterbalances the fit-to-cohort prioritization inherent in classic genetic-only approaches, which have been plagued by poor reproducibility across cohorts. We have now extended our previous work to include more datasets of GWAS, and more recent evidence from other lines of work. In essence our analysis is the most comprehensive integration of genetics and functional genomics to date in the field of bipolar disorder. Biological pathway analyses identified top canonical pathways, and epistatic interaction testing inside these pathways has identified genes that merit future follow-up as direct interactors (intra-pathway epistasis, INPEP). Moreover, we have put together a panel of best *P*-value single nucleotide polymorphisms (SNPs), based on the top candidate genes we identified. We have developed a genetic risk prediction score (GRPS) based on our panel, and demonstrate how in two independent test cohorts the GRPS differentiates between subjects with bipolar disorder and normal controls, in both European-American and African-American populations. Lastly, we describe a prototype of how such testing could be used to categorize disease risk in individuals and aid personalized medicine approaches, in psychiatry and beyond.

© 2010 Wiley-Liss, Inc.

Key words: convergent functional genomics; pathways; epistasis; genetic risk; prediction; bipolar disorder

INTRODUCTION

As part of a convergent functional genomics (CFG) strategy, expanding upon our earlier work [Le-Niculescu et al., 2009b], we

How to Cite this Article:

Patel SD, Le-Niculescu H, Koller DL, Green SD, Lahiri DK, McMahon F, Nurnberger JJ, Niculescu AB. 2010. Coming to Grips With Complex Disorders: Genetic Risk Prediction in Bipolar Disorder Using Panels of Genes Identified Through Convergent Functional Genomics.

Am J Med Genet Part B 9999:1–28.

set out to comprehensively identify candidate genes for bipolar disorder, integrating available evidence in the field to date. We have used data from four publicly available genome-wide association studies (GWAS) datasets for bipolar disorder [WTCC, 2007; Baum et al., 2008; Sklar et al., 2008]. We integrated those data with gene expression data—human postmortem brain gene expression data and human blood gene expression data published by others or us, as well as with relevant animal model brain and blood gene expression data generated by our group

Additional Supporting Information may be found in the online version of this article and at www.neurophenomics.info

ABN is a scientific co-founder of Mindscape Diagnostics.

*Correspondence to:

Prof. A.B. Niculescu, M.D., Ph.D., Assistant Professor of Psychiatry and Medical Neuroscience, Indiana University School of Medicine, Staff Psychiatrist, Indianapolis VA Medical Center, Director, INBRAIN and Laboratory of Neurophenomics, Institute of Psychiatric Research, 791 Union Drive, Indianapolis, IN 46202-4887. E-mail: anicules@iupui.edu
Published online 00 month 2010 in Wiley InterScience
(www.interscience.wiley.com)

DOI 10.1002/ajmg.b.31087

[Niculescu et al., 2000a; Ogden et al., 2004; Le-Niculescu et al., 2007a, 2007b, 2008b]. In addition, we have integrated as part of this comprehensive approach other genetic data-published human genetic (linkage or association) data for bipolar and related disorders to date, and relevant mouse genetic (QTL or transgenic) data (Fig. 1).

Once the genes involved in a disorder are identified, and prioritized for likelihood of involvement, then an obvious next step is developing a way of applying that knowledge to genetic testing of individuals to determine risk for the disorder. Based on our comprehensive identification of top candidate genes described in this paper, we have chosen the best SNPs in those genes by their *P* values in the GWAS datasets used, and assembled a genetic risk prediction (GRP) panel out of those SNPs. We then developed a genetic risk prediction score (GRPS) for bipolar disorder based on the presence or absence of the alleles of the SNPs associated with the illness, and tested the GRPS in an independent study (GAIN-BP) [Smith et al., 2009] for which we had both genotypic and clinical data available, comparing the bipolar subjects to demographically matched normal controls. Our results show that a relatively small size panel of genes identified by CFG analysis can differentiate very well between bipolar disorder subjects and controls at a population level, although at an individual level the margin is razor thin. The latter point suggests that the cumulative combinatorics of common variants plays a major role in risk for illness. Overall, our work sheds light on the genetic architecture and pathophysiology of bipolar disorder. In particular, it has implications for genetic testing to assess risk for illness before the illness manifests itself clinically.

METHODS

Genome-Wide Association Studies (GWAS) Data for Bipolar Disorder

Four bipolar GWAS were used for the expanded CFG discovery analysis. The GWAS data for the bipolar study from the Wellcome Trust Consortium (WTCC) [2007] is available at http://www.wtccc.org.uk/info/access_to_data_samples.shtml. The GWAS data from NIMH and German studies [Baum et al., 2008] are available at http://mapgenetics.nimh.nih.gov/bp_pooling. The GWAS data from the STEP-BD study is available at <http://pngu.mgh.harvard.edu/~purcell/bpwgas> [Sklar et al., 2008].

One independent study, GAIN-BP [Smith et al., 2009], was used for testing the results of the discovery analyses. The GWAS data for GAIN-BP used for analyses described in this manuscript was obtained from the database of Genotype and Phenotype (dbGaP) found at www.ncbi.nlm.nih.gov through PHS project number 000017, data request numbers 2575-2, 2574-2, and 2573-2 provided to John I. Nurnberger, Jr.

The software package PLINK (<http://pngu.mgh.harvard.edu/~purcell>) was used to extract individual genotype information for each subject from the GAIN-BP GWAS data files. We used European Americans (EA), and separately, African American (AA), bipolar subjects and controls. Out of 1001 EA bipolar subjects in GAIN-BP, we used for our GRPS testing analyses only 407, from wave 5 of the NIMH Bipolar Genetics Consortium collection, to avoid any individual overlap (16%) or even pedigree overlap (57%) with probands from waves 1 to 4 that were also used in the NIMH [Baum et al., 2008] study mentioned above. We also used 317 AA bipolar subjects from wave 5 of the NIMH Bipolar Genetics Consortium collection. Controls numbered 1034 for EA, and 671 for AA.

As a caveat, there was overlap in the control subjects within two of four discovery datasets (NIMH and STEP-BD), and between these two datasets and one of the datasets (GAIN-BP EA) used to test our results. However, as described below, multiple other studies and lines of evidence, human and animal model, are integrated in the CFG prioritization approach (Fig. 1), which minimizes the relative contribution and impact of individual studies and the controls overlap in the discovery dataset. More importantly, there is no overlap at the bipolar subject level within discovery datasets, and between discovery datasets and the test dataset. This ensures that there is at least a degree of independence within discovery cohorts, and between discovery and test cohorts. Finally, the fact that the GRPS differentiates as well or better in the completely independent GAIN-BP AA cohort provides strong reassurance and confirmatory evidence for the method.

SNPs with a nominal genotypic *P*-value <0.05 were selected for our analysis. No Bonferroni correction was performed.

Gene Identification

To identify the genes that correspond to the selected SNPs, the lists of SNPs from the GWAS was uploaded to the CHIP Bioinformatics Tools website (<http://snpper.chip.org>). In the cases where a SNP mapped to a region close to multiple genes, we selected all the genes




FIG. 1. Convergent functional genomics. Integration of multiple independent lines of evidence. The maximal possible score from GWAS data [6 pt] is equally weighed with the maximal possible score from other lines of evidence [other human and animal model gene expression and genetic data] [6 pt].

that were provided by SNPper. SNPs for which no gene was identified were not included in our subsequent analysis.

Human Postmortem Brain Gene Expression

Information about our candidate genes was obtained using GeneCards (<http://www.genecards.org>), the Online Mendelian Inheritance of Man database (<http://ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM>), as well as database searches using PubMed (<http://ncbi.nlm.nih.gov/PubMed>) and various combinations of keywords (gene name, bipolar, depression, human, postmortem, brain).

Human Genetic (Linkage, Association) Convergence

To designate convergence for a particular gene, the gene had to map within 10 cM [see Niculescu et al., 2000b for detailed discussion] of a microsatellite marker for which at least one published study showed evidence for linkage for bipolar disorder or depression, or a positive association study for the gene itself was reported in the literature. The University of Southampton's sequence-based integrated map of the human genome (The Genetic Epidemiological Group, Human Genetics Division, University of Southampton: http://cedar.genetics.soton.ac.uk/public_html/) was used to obtain cM locations for both genes and markers. The sex-averaged cM value was calculated and used to determine convergence to a particular marker. For markers that were not present in the Southampton database, the Marshfield database (Center for Medical Genetics, Marshfield, WI, USA: <http://research.marshfieldclinic.org/genetics>) was used with the NCBI Map Viewer web-site to evaluate linkage convergence.

We have established in the lab manually curated databases of all the published human postmortem brain and human genetic literature to date on bipolar and related disorders [Niculescu and Le-Niculescu, 2010]. These large databases have been used in our CFG cross-validation analyses.

Human Blood Gene Expression Data

For human blood gene expression evidence, we have used previously generated data from our group [Le-Niculescu et al., 2009a], as well as published data from the literature.

Animal Model Brain and Blood Gene Expression Data

For animal model brain and blood gene expression evidence, we have used previously generated data from two different animal models for bipolar disorder developed by our group, one pharmacogenomic and one transgenic [Niculescu et al., 2000a; Le-Niculescu et al., 2007b, 2008c].

Mouse Genetic (QTL, Transgenic) Convergence

To search for mouse genetic evidence—quantitative trait loci (QTL) or transgenic—for our candidate genes, we utilized the MGI_3.54—Mouse Genome Informatics (Jackson Laboratory, Bar Harbor, Maine) and used the search menu for mouse phenotypes and mouse models of human disease/abnormal behaviors, using the

following sub-categories: abnormal emotion/affect behavior and abnormal sleep pattern/circadian rhythm. To designate convergence for a particular gene, the gene had to map within 10 cM of a QTL marker for the abnormal behavior, or a transgenic mouse of the gene itself displayed that behavior.

Convergent Functional Genomics (CFG) Analysis Scoring

We used two nominal *P*-value thresholds for scoring genes in the CFG analysis (see below) a lower stringency threshold ($P < 0.05$), and a higher stringency threshold ($P < 0.001$). Genes from each GWAS data that had at least one SNP with *P*-value of < 0.05 received 1 point; those that had at least one SNP with *P*-value of < 0.001 received 1.5 points. All other cross-validating lines of evidence (other human data, animal model data) received a maximum of 1 point each (for human genetic data: 0.5 points if it is linkage, 1 point if it is association; for mouse genetic data, 0.5 points if it is QTL, 1 point if it is transgenic; for human and mouse gene expression data, 1 point each for brain or blood data, 0.5 points if it is from cells in culture/cell lines). Thus the maximum possible CFG score for each gene is 12 ($6 = 4 \times 1.5$ points from the four GWAS, and 6 points from the other lines of evidence). As we are interested in discovering signal in GWAS, we weighted data from GWAS more heavily, bringing the data from this one methodological approach on par with the data from all the other methodological approaches combined. It has not escaped our attention that other ways of weighing the scores of line of evidence may give slightly different results in terms of prioritization, if not in terms of the list of genes per se. Nevertheless, we feel this simple scoring system provides a good separation of genes based on our focus on identifying signal in the GWAS.

Pathway Analysis

Ingenuity 8.0 (Ingenuity Systems, Redwood City, CA) was employed to analyze the molecular networks, biological functions and canonical pathways of the top candidate genes resulting from our CFG analysis. The Ingenuity program generated the *P* values assigned to the different pathways (Table I).

Epistasis Testing

The GAIN-BP case and control data were employed to test for epistatic interactions among SNPs in genes from the GRPS panel having a role in one or more of the top canonical biological pathways from our pathway analysis. These pathways, and the genes comprising each that were considered, are listed in Table II. Within each pathway, SNP \times SNP allelic epistasis was tested for each distinct pair of SNPs using the PLINK software package.

Genetic Risk Prediction Panel and Scoring

Out of our analysis, a panel of top genes prioritized by CFG scoring (Fig. 1) can be chosen. We developed a genetic risk prediction (GRP) panel, based on a list of top genes from Table III ($n = 56$, all the genes that had a CFG score better than 6, that is, $>$ than 50% of the maximum possible CFG score of 12). All the SNPs for these

TABLE I. Top Candidate Genes for Bipolar Disorder Identified by Convergent Functional Genomics (CFG) of Genome-Wide Association Studies (GWAS) Data and Replication of Findings in an Independent Study (GAIN-BP)

Entrez ID	Gene symbol/name	GWAS WTCC [2007] P-value	GWAS NIMH [Baum et al., 2008], P-value	GWAS German [Baum et al., 2008], P-value	GWAS step-BD [Sklar et al., 2008], P-value	Mouse models brain evidence [Ogden et al., 2004; Le-Niculescu et al., 2008c] PFC (D) Cat IV-meth	Mouse models blood evidence [Le-Niculescu et al., 2008b] PFC (D) Cat IV-meth	Human genetic evidence (linkage or association) 11p15.2	Human postmortem brain evidence (BP) [Nakatani et al., 2006]	Human blood/other peripheral tissue evidence	CFG score	GAIN-BP P-value
406	ARNTL (Argy hydrocarbon receptor nuclear translocator-like)	7.71E-04, rs4757141	3.84E-02, rs4757138	3.72E-02, rs3816360	2.56E-02, rs11022781	Abnormal sleep pattern/circadian rhythm (TG)					8.5	4.87E-02
4155	MBP (Myelin basic protein)	8.30E-03, rs470549	8.19E-04, rs470821	1.17E-03, rs12967023		NST PFC (D) ST PFC (D) ST AMY (I)	BP Blood (I) Cat IV-meth	18q23 (assoc)	D(BP) [Kachev et al., 2003]		8.5	3.14E-03
627	BDNF (Brain derived neurotrophic factor)	1.05E-02, rs16917234	3.76E-02, rs925946	1.91E-03, rs12291063		Abnormal emotion/affect behavior (TG)	PFC (D) Cat IV-METH	11p14.1 (assoc)	D(BP) [Knable et BP (D) al., 2004; Torrey et al., 2005; Pillai, 2008]	Mood (I) [Le-Niculescu et al., 2009a] D (female BP) [Chambers and Perrone-Bizzozero, 2004]	8.0	4.19E-02

BP [Mansour et al., 2006; Nievergelt et al., 2006; Partonen et al., 2007]

(assoc)

BP [Potash et al., 2008] (assoc)

BP [Freimer et al., 1996; Schulze et al., 2003; Mazziade et al., 2005]

BP [Neves-Pereira et al., 2002; Sklar et al., 2002; Liu et al., 2008] (assoc)

BP [McInnes et al., 1996; Detera-Wadleigh et al., 1999]

Depression [Aguilera et al., 2009] (assoc)
MDD [Licinio et al., 2009] (assoc)

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS			GWAS step-BD [Sklar et al., 2008]. P-value	Mouse genetic evidence (QTL, TG)	Mouse models brain evidence		Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	CFG score	GAIN-BP P-value
		NIMH [Baum et al., 2008], P-value	German [Baum et al., 2008], P-value	WTCC [2007] P-value			[Ogden et al., 2004; Le-Niculescu et al., 2008c]	[Le-Niculescu et al., 2008b]					
3084	NRG1 (Neuregulin 1)	2.19E-03, rs327380	4.51E-03, rs6468095	1.07E-05, rs7821190	8.81E-04, rs2466085		Le-Niculescu et al., 2008c	[Le-Niculescu et al., 2008b]	8p12	I [BP] [Tkachev BP (I) et al., 2003]	[Begemann et al., 2008]	8.0	1.16E-02
6096	RORB (RAR-related orphan receptor beta)	5.88E-04, rs1327837	1.95E-02, rs1359073	1.29E-02, rs10869435	8.99E-04, rs10869435	Abnormal emotion/affect behavior [TG]	ST AMY (I) ST PFC (D)		9q21.13	BP [Green et al., 2005; Waliss-Bass et al., 2006; Thomson et al., 2007; Georgieva et al., 2008] (assoc)	D (unipolar depression) [Bertram et al., 2007]	8.0	9.38E-03
27185	DISC1 (Disrupted in schizophrenia 1)	2.99E-03, rs9431714	6.08E-03, rs7534681	1.31E-02, rs6671423	8.61E-03, rs821577	Abnormal emotion/affect behavior [TG]			1q42.2	BP [Macgregoret al., 2004] BP [McGrath et al., 2009] (assoc)	MDD [Sawamura BP(D) et al., 2005]	8.0	2.73E-02
54715	A2BP1 (Ataxin-2-binding protein 1)	4.23E-04, rs1818290	1.59E-04, rs11077135	3.42E-05, rs8046170	4.18E-03, rs7187986		VT (D) Cat III-VPA		16p13.2	BP [Hodgkinson et al., 2004; Thomson et al., 2005; Maeda et al., 2006; Perlis et al., 2008] (assoc)	MDD [Schosser et al., 2009] (assoc)	7.5	1.11E-02
										BP [Baum et al., 2008; Johnson et al., 2009] (assoc)	BP [Ewald et al., 2002]		

(Continued)

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS		GWAS WTCC [2007]	GWAS NIMH [Baum et al., 2008], P-value	GWAS German [Baum et al., 2008], P-value	GWAS step-BD [Sklar et al., 2008], P-value	Mouse models brain evidence [Ogden et al., 2004; Le-Niculescu et al., 2008c]	Mouse models blood evidence [Le-Niculescu et al., 2008b]	Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	CFG score	GAIN-BP P-value
		P-value	P-value											
216	ALDH1A1 (Aldehyde dehydrogenase family 1, subfamily A1)	1.29E-02, rs348478	1.58E-04, rs348458	1.29E-02, rs348478	1.58E-04, rs348458	3.34E-02, rs7873724		Abnormal sleep pattern/circadian rhythm (QTL)	BP blood (D) Cat IV-Meth	9q21.13	I (BP) [Pennington et al., 2008]		7.5	3.08E-02
2890	GRIA1 (Glutamate receptor, ionotropic, AMPA 1)	1.47E-02, rs17096210	6.55E-03, rs4958667	1.47E-02, rs17096210	6.55E-03, rs4958667	9.19E-03, rs7719292	6.84E-03, rs1461232	Abnormal emotion/affect behavior (QTL)	VT (D) Cat IV-Meth	5q33.2	I (BP), (MDD) [Choudary et al., 2005]		7.5	2.48E-04
4842	NOS1 (Nitric oxide synthase 1), neuronal	1.72E-02, rs1607817	3.73E-02, rs2271987	1.72E-02, rs1607817	3.73E-02, rs2271987	4.56E-02, rs12811583	2.12E-02, rs10850803	Abnormal emotion/affect behavior abnormal sleep pattern/circadian rhythm (QTL)	NST AMY (D)	12q24.22	I/D (BP) [Benes et al., 2006]		7.5	1.13E-02
773	CACNA1A (Calcium channel, voltage-dependent, P/Q type, alpha 1A) subunit	2.99E-02, rs1777941	2.12E-02, rs10421810	2.99E-02, rs1777941	2.12E-02, rs10421810	7.04E-04, rs10421810	2.70E-03, rs16016	Abnormal emotion/affect behavior (QTL)		19p13.13	D (BP) [Iwamoto et al., 2004]		7.0	5.67E-03
10659	CUGBP2 (CUG triplet repeat, RNA binding protein 2)	2.84E-05, rs682970	3.38E-03, rs932918	2.84E-05, rs682970	3.38E-03, rs932918	2.66E-02, rs2378991	9.51E-03, rs1990	ST PFC (I)		MDD [Zubenko et al., 2003] 10p14	BP (I) [Matigian et al., 2007]		7.0	1.59E-02
										MDD [Zubenko et al., 2003]				

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS		GWAS WTCC [2007] P-value	GWAS		GWAS step-BD [Sklar et al., 2008], P-value	Mouse models brain evidence [Ogden et al., 2004; Le-Niculescu et al., 2008c]	Human genetic evidence [linkage or association]	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	CFG score	P-value
		NIMH [Baum et al., 2008], P-value	German [Baum et al., 2008], P-value										
1826	DSCAM (Down syndrome cell adhesion molecule) like 1	2.72E-04, rs8129283	8.11E-04, rs7228073	1.39E-03, rs455304	1.51E-02, rs2837504	21q22.2	I (BP) [Amano et al., 2008]	Human genetic evidence [linkage or association]	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	7.0	1.66E-03	
2913	GRM3 (Glutamate receptor, metabotropic 3)	3.18E-03, rs10226401	7.36E-03, rs2237552	3.43E-02, rs6955917	2.22E-04, rs2237554	ST AMY (I)	BP [Amano et al., 2008] (assoc) 7q21.12	D (MDD/suicide) [Klempan et al., 2009]	D (BP) [Choudhary et al., 2005; Etain et al., 2005]	D (BP) [Nakatani et al., 2006; Vawter et al., 2006]	7.0	5.69E-03	
2932	GSK3B (Glycogen synthase kinase 3 beta)	1.62E-02, rs17810235	6.72E-03, rs6438552	9.82E-03, rs17811013	Abnormal emotion/affect behavior (TG)	CP (D) Cat IV-VPA ST PFC (D) ST AMY (I) PFC (D) Cat IV-METH	BP [Lambert et al., 2005; Etain et al., 2006]	3q13.33	D (BP) [Nakatani et al., 2006; Vawter et al., 2006]	I (MDD) [Vawter et al., 2006]	7.0		
3356	HTR2A (5-Hydroxytryptamine (serotonin) receptor 2A)	4.52E-02, rs972979	1.65E-03, rs17288723	1.86E-02, rs2025296	5.60E-03, rs977003	Abnormal emotion/affect behavior, abnormal sleep pattern/circadian rhythm (TG)	13q14.2	D (BP) [Knable et al., 2004; Torrey et al., 2005]	D (BP) [Knable et al., 2004; Torrey et al., 2005]	I (Suicide) [Arranz et al., 1997; Lin et al., 2003; McAuley et al., 2009] (assoc)	7.0	3.19E-03	

(Continued)

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS			Mouse genetic evidence (QTL, TG)	Mouse models brain evidence		Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/peripheral tissue evidence	CFG score	GAIN-BP P-value
		WTCC [2007] P-value	NIMH [Baum et al., 2008], P-value	German [Baum et al., 2008], P-value		step-BD [Sklar et al., 2008], P-value	Le-Niculescu et al., 2008c					
3775	KCNK1 (Potassium channel, subfamily K, member 1)	1.89E-02, rs3843250	7.60E-03, rs4649240	3.47E-04 , rs701209	4.38E-02, rs4649343	Ogden et al., 2004; Le-Niculescu et al., 2008c	Human genetic evidence (linkage or association) Major affective disorders [Bonnier et al., 2002] (assoc) Mood disorders [Brezo et al., 2009] (assoc) Response to antidepressants (SSRI) [Uher et al., 2009] (assoc) 1q42.2 D (BP) [Jurata et BP (I) [Matigian et al., 2004]	Human postmortem brain evidence	Human blood/peripheral tissue evidence	7.0		
11278	KLF12 (Kruppel-like factor 12)	2.76E-03, rs4885151	6.77E-04 , rs9600160	1.68E-04 , rs9543443	Abnormal emotion/affect behavior	ST AMY (I) ST PFC (D)	BP [Curtis et al., 2003; Macgregor et al., 2004] 13q22.1	Mood (D) [Le-Niculescu et al., 2009a]		7.0	4.93E-04	
10150	MBNL2 (Muscleblind-like 2)	2.94E-03, rs6491345	4.64E-02, rs7318623	4.02E-04 , rs9584552	1.61E-02, rs16953952	AMY (D) Cat III-VPA	BP [Potash et al., 2003] 13q32.1			7.0	4.08E-02	
89797	NAV2 (Neuron navigator 2)	4.16E-03, rs2119981	5.77E-04 , rs1372797	2.04E-03, rs2218329	1.87E-03, rs10500860	Abnormal emotion/affect behavior (TG)	BP [Keisoe et al., 2001] 11p15.1 D, bipolar suicide [Kim et al., 2007]			7.0	1.73E-03	
							BP [Detera-Wadleigh et al., 1999]					

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS			Mouse genetic evidence (QTL, TG)	Mouse models evidence [Ogden et al., 2004; Le-Niculescu et al., 2008c]	Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	CFG score	GAIN-BP P-value
		GWAS NIMH [Baum et al., 2008], P-value	GWAS German [Baum et al., 2008], P-value	GWAS step-BD [Sklar et al., 2008], P-value							
4684	NCAM1 (Neural cell adhesion molecule 1)	GWAS WTCC [2007] P-value 2.77E-02, rs11214501	GWAS NIMH [Baum et al., 2008], P-value 2.61E-02, rs586903	GWAS German [Baum et al., 2008], P-value 8.62E-03, rs12279261	Abnormal emotion/affect behavior (TG)	[Le-Niculescu et al., 2008b]	11q23.1	D [BP] [Atz et al., 2007], MDD [D] [Tochigi et al., 2008]	peripheral blood cells [Wakabayashi et al., 2008]	7.0	
2908	NR3C1 (Nuclear receptor subfamily 3, group C, member 1) (glucocorticoid receptor)	4.03E-03, rs17209251	3.71E-02, rs10482672	2.96E-02, rs10482672	Abnormal emotion/affect behavior (TG)	[Le-Niculescu et al., 2008b]	5q31.3	D [BP] [Knable et al., 2004; Torrey et al., 2005]		7.0	7.54E-03
4988	OPRM1 (Opioid receptor, mu 1)	7.82E-04, rs2010884	7.31E-03, rs650825	1.90E-03, rs7745499	Abnormal emotion/affect behavior (TG)		6q25.2	I [BP] [Ryan et al., 2006]		7.0	7.57E-03
5101	PCDH9 (Protocadherin 9)	9.77E-03, rs17082149	1.19E-03, rs9317626	4.80E-04, rs7986387	Abnormal emotion/affect behavior, abnormal sleep pattern/circadian rhythm (QTL)	NST AMY (I)	13q21.32	D [MDD/suicide] [Klempner et al., 2009]		7.0	4.92E-03

(Continued)

TABLE 1. (Continued)

Entrez ID	Gene symbol/name	GWAS WTCC [2007]		GWAS NIMH [Baum et al., 2008]		GWAS German [Baum et al., 2008]		GWAS step-BD [Sklar et al., 2008]		Mouse models brain evidence [Ogden et al., 2004]; Le-Niculescu et al., 2008c]	Mouse models blood evidence [Le-Niculescu et al., 2008b]	Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	CFG score	GAIN-BP P-value
		P-value	rs	P-value	rs	P-value	rs	P-value	rs							
5142	PDE4B (Phosphodiesterase 4B), cAMP-specific (phosphodiesterase E4 dunce homolog, drosophila)	6.02E-03, rs6588190	1.60E-03, rs539322	1.41E-02, rs12021574	4.42E-02, rs17417507	1.20E-02, rs6748375	2.48E-02, rs4557033	Abnormal emotion/affect behavior (TG)	CP (I) Cat IV-Meth	2p21	D (BP) [Torrey et al., 2005]	D (BP) [Fatemi BP (I) [Padmos et al., 2008]	D (BP) [Torrey et al., 2005]	MDD (I), leukocyte [Numata et al., 2009]	7.0	1.46E-03
5581	PRKCE (Protein kinase C, epsilon)	4.59E-03, rs2711293	2.37E-04 , rs2595221	1.20E-02, rs6748375	2.48E-02, rs4557033	1.90E-02, rs6977749	1.01E-02, rs17169022	Abnormal emotion/affect behavior (TG)	CP (I) Cat IV-Meth	2p21	D (BP) [Torrey et al., 2005]	D (BP) [Fatemi BP (I) [Padmos et al., 2008]	D (BP) [Torrey et al., 2005]	MDD (I), leukocyte [Numata et al., 2009]	7.0	1.46E-03
5764	PTN (Pleiotrophin) (heparin binding growth factor 8, neurite growth-promoting factor 1)	2.85E-02, rs6977819	1.90E-02, rs6977749	4.56E-03, rs320682	8.78E-04 , rs17169022	1.90E-02, rs6977749	1.01E-02, rs17169022	Abnormal emotion/affect behavior (TG)	CP (I) Cat IV-Meth	7q33	I (MDD) [Tochigi et al., 2008]	I (MDD) [Tochigi et al., 2008]	I (MDD) [Tochigi et al., 2008]	MDD (I), leukocyte [Numata et al., 2009]	7.0	1.03E-03
5797	PTPM (Protein tyrosine phosphatase, receptor type, M)	1.74E-02, rs727951	1.10E-02, rs3786367	2.41E-04 , rs16952620	1.01E-02, rs4121619	6.09E-03, rs2670955	8.07E-03, rs744776	Abnormal emotion/affect behavior (TG)	CP (I) Cat IV-VPA	18p11.23	I (BP) [Nakatani Mood et al., 2006]	I (BP) [Nakatani Mood et al., 2006]	I (BP) [Nakatani Mood et al., 2006]	Mood (D) [Le-Niculescu et al., 2009a]	7.0	1.14E-03
6263	RYR3 (Ryanodine receptor 3)	1.21E-03, rs16957945	2.89E-04 , rs2596205	6.09E-03, rs2670955	8.07E-03, rs744776	6.09E-03, rs2670955	8.07E-03, rs744776	Abnormal emotion/affect behavior (TG)	CP (I) Cat IV-VPA	15q13.3	BP [Segurado et al., 2003]	BP [Segurado et al., 2003]	BP [Segurado et al., 2003]	BP [Segurado et al., 2003]	7.0	1.11E-03
9522	SCAMP1 (Secretory carrier membrane protein 1)	1.71E-02, rs1019803	1.31E-02, rs1968382	2.46E-03, rs16875382	2.25E-02, rs16875428	2.46E-03, rs16875382	2.25E-02, rs16875428	Abnormal emotion/affect behavior (OTL)	ST PFC (D)	5q14.1	Depression [Levinson et al., 2007]	Depression [Levinson et al., 2007]	Depression [Levinson et al., 2007]	Mood (D) [Le-Niculescu et al., 2009a]	7.0	4.14E-02
287	ANK2 (Ankyrin 2), neuronal	4.77E-04 , rs17445459	1.34E-02, rs17590593	8.90E-03, rs10516593	5.18E-03, rs1351998	8.90E-03, rs10516593	5.18E-03, rs1351998	Abnormal emotion/affect behavior (OTL)	ST PFC (I)	4q25	BP [Lambert et al., 2005]	BP [Lambert et al., 2005]	BP [Lambert et al., 2005]	BP [Lambert et al., 2005]	6.5	9.42E-05

TABLE 1. (Continued)

Entrez ID	Gene symbol/name	GWAS			Mouse models evidence [Ogden et al., 2004; Le-Niculescu et al., 2008c]	Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	CFG score	GAIN-BP P-value
		WTCC [2007] P-value	NIMH [Baum et al., 2008], P-value	German [Baum et al., 2008], P-value						
351	APP (Amyloid beta(A4) precursor protein)	3.37E-02, rs3991	9.86E-03, rs2829984	7.81E-03, rs3787620	1.04E-02, rs2830048	Abnormal emotion/affect behavior, abnormal sleep pattern/circadian rhythm (TG)	21q21.3 I (BP) [Jurata et al., 2004]		6.5	
6310	ATXN1 (Ataxin 1)	1.11E-03, rs9370893	5.55E-03, rs2237198	6.58E-03, rs12198838	3.96E-04, rs909786			Mood (I)	6.5	1.12E-03
815	CAMK2A (Calcium/calmodulin-dependent protein kinase II alpha)	1.76E-02, rs10515639	3.62E-02, rs3797617	3.62E-02, rs3797617	2.30E-02, rs4958469	Abnormal emotion/affect behavior, abnormal sleep pattern/circadian rhythm (TG)	5q32 D (BP) [Xing et al., 2002], I (MDD) [Novak et al., 2006; Tochigi et al., 2008]		6.5	
960	CD44 (CD44 antigen)	3.48E-02, rs16927100	3.94E-03, rs353615	1.06E-02, rs7115768				BP [Morissette et al., 1999] 6p22.3		
1012	CDH13 (Cadherin 13)	5.89E-03, rs1862682	2.50E-03, rs7198252	9.08E-04, rs931408	8.01E-03, rs7197423	Abnormal emotion/affect behavior (OTL)	BP [Sklar et al., 2004; Etain et al., 2006] 11p13	BP blood (D) Cat IV-Meth	6.5	4.01E-04
1387	CREBBP (CREB binding protein)	5.02E-03, rs130036	1.39E-03, rs13332076	3.64E-03, rs129963	1.91E-02, rs11644593	Abnormal emotion/affect behavior (TG)	BP [Etain et al., 2006] 16p13.3		6.5	1.37E-03
1612	DAPK1 (Death-associated protein kinase 1)	4.02E-02, rs11141909	5.97E-05, rs10868644	4.04E-02, rs3124236	1.56E-03, rs3124236	Abnormal emotion/affect behavior (OTL)	BP [Ewald et al., 2002] 9q21.33		6.5	5.20E-03

(Continued)

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS WTCC [2007]		GWAS NIMH [Baum et al., 2008]		GWAS German [Baum et al., 2008]		GWAS step-BD [Sklar et al., 2008], P-value		Mouse genetic evidence (QTL, TG)	Mouse models brain evidence [Ogden et al., 2004; Le-Niculescu et al., 2008c]	Mouse models blood evidence [Le-Niculescu et al., 2008b]	Human genetic evidence (linkage or association) BP [Segurado et al., 2003]	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	CFG score	GAIN-BP P-value
		P-value	rs	P-value	rs	P-value	rs	P-value	rs								
9201	DCLK1 [Doublecortin-like kinase 1]	1.20E-02	rs740474	2.36E-03	rs11954658	5.27E-03	rs397327	2.59E-02	rs3792896	BP (D) Cat IV-VPA	DBP NST (D)		BP [Etain et al., 2006] 3p13			6.5	
1729	DIAPH1 [Diaphanous (Drosophila, homolog 1)]	2.62E-02, rs740474	rs740474	4.70E-02, rs11954658	rs11954658	3.38E-03, rs397327	rs397327	7.63E-03, rs3792896	rs3792896	CP (I) Cat III-VPA			BP, SZ [Maziade et al., 2005] 5q31.3		BP (I) [Matigian et al., 2007]	6.5	
27086	FOXP1 [Forkhead box P1]	4.80E-03, rs950443	rs950443	9.66E-04 , rs7640237	rs7640237	5.33E-03, rs3846031	rs3846031	1.58E-04 , rs17718783	rs17718783	NST AMY (D) ST PFC (D)			BP [Etain et al., 2006] 3p13			6.5	2.97E-03
2770	GNAI1 [Guanine nucleotide binding protein, alpha inhibiting 1]	4.98E-03, rs10429156	rs10429156	7.55E-03, rs6973616	rs6973616	1.55E-02, rs916905	rs916905	3.71E-02, rs2523189	rs2523189	ST PFC (D)			BP [McInnes et al., 1996; Etain et al., 2006] 7q21.11	D (BP) [Jurata et al., 2004]		6.5	4.32E-02
2897	GRIK1 [Glutamate receptor, ionotropic, kainate 1]	5.39E-04 , rs2154490	rs2154490	2.79E-03, rs2832476	rs2832476	3.36E-02, rs464982	rs464982	4.47E-02, rs467155	rs467155	Abnormal emotion/affect behavior (QTL)			BP [Lambert et al., 2005] 21q21.3	D (BP) [Iwamoto et al., 2004; Choudary et al., 2005; Nakatani et al., 2006]		6.5	5.91E-03
2939	GSTA2 [Glutathione S-transferase, alpha 2] [Yc2]	1.14E-03, rs2207950	rs2207950	1.93E-03, rs2608632	rs2608632	1.89E-03, rs2224198	rs2224198	1.52E-02, rs2749010	rs2749010		BP (D) Cat III-Meth		BP [Detera-Wadleigh et al., 1999; Morissette et al., 1999] 6p12.2	I (BP) [Benes et al., 2006]		6.5	
3751	KCND2 [Potassium voltage-gated channel, Shal-related family, member 2]	5.78E-03, rs10156125	rs10156125	4.08E-03, rs12538990	rs12538990	5.24E-05 , rs10268591	rs10268591	3.86E-02, rs2191736	rs2191736	ST PFC (D)			BP [Lambert et al., 2005] 7q31.31			6.5	
													BP [Etain et al., 2006]				

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS		GWAS		GWAS		GWAS		Mouse models brain evidence		Human genetic evidence		Human postmortem brain evidence		Human blood/peripheral tissue evidence		CFG score	GAIN-BP P-value
		WTCC [2007] P-value	NIMH [Baum et al., 2008], P-value	Baum German [Baum et al., 2008], P-value	Sklar et al., [Sklar et al., 2008], P-value	step-BD [Sklar et al., 2008], P-value	Mouse genetic evidence (QTL, TG)	Mouse models blood evidence [Le-Niculescu et al., 2008c]	Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/peripheral tissue evidence	CFG score	GAIN-BP P-value						
4008	LM07 [LIM domain only ?]	6.62E-05, rs9530460	1.11E-02, rs9593132	8.17E-03, rs1570554	6.59E-03, rs9530460	Abnormal emotion/affect behavior, abnormal sleep pattern/circadian rhythm (QTL)	BP [Potash et al., 2003] 2p25.3	13q22.2	Human postmortem brain evidence	Human blood/peripheral tissue evidence	6.5	2.26E-02	Anti-depressant (D) lymphocytes [Kaiman et al., 2005]						
23040	MYT1L (Myelin transcription factor 1-like)	2.25E-04, rs1991773	1.31E-02, rs1421614	1.25E-02, rs10519486	1.65E-02, rs17039396	Abnormal sleep pattern/circadian rhythm (QTL)	BP [Potash et al., 2003] 2p25.3	ST PFC (D)	Human genetic evidence (linkage or association)	Human postmortem brain evidence	6.5	5.83E-03							
4720	NDUFS2 (NADH dehydrogenase (ubiquinone) Fe-S protein 2), 49 kDa [NADH-coenzyme Q reductase])	4.27E-02, rs5085	1.08E-02, rs11421	4.67E-02, rs11421	3.61E-02, rs5085	BP (I) Cat III-VPA	BP [Detera-Wadleigh et al., 1999] 1q23	BP (I) Cat III-VPA	Human genetic evidence (linkage or association)	Human postmortem brain evidence	6.5		BP (D) [Middleton et al., 2005]						
4897	NRCAM (Neuronal cell adhesion molecule)	1.63E-03, rs13227836	5.94E-04, rs3763461	8.60E-04, rs1548949	4.35E-02, rs11974528	Abnormal sleep pattern/circadian rhythm (QTL)	BP [Fallin et al., 2004] 7q31.1	NST AMY (I)	Human genetic evidence (linkage or association)	Human postmortem brain evidence	6.5	1.16E-03							
10846	PDE10A (Phosphodiesterase 10A)	1.50E-02, rs2983506	9.64E-03, rs12525763	1.50E-03, rs454165	3.40E-03, rs2983521	Abnormal emotion/affect behavior (TG)	6q27	NST AMY (D) ST PFC (D)	Human genetic evidence (linkage or association)	Human postmortem brain evidence	6.5	2.16E-02							
11122	PTPR1 (Protein tyrosine phosphatase, receptor type, T)	6.27E-03, rs6030385	3.45E-03, rs2425478	1.12E-02, rs1016071	4.67E-03, rs1883842	ST AMY (I)	BP [Cheng et al., 2006] 20q12	I (MDD/suicid) [Sequeira et al., 2007]	Human genetic evidence (linkage or association)	Human postmortem brain evidence	6.5	1.20E-03	I (MDD) [Aston et al., 2005]						

(Continued)

TABLE I. (Continued)

Entrez ID	Gene symbol/name	GWAS		GWAS		GWAS		GWAS		Mouse models brain evidence		Human genetic evidence		Human postmortem brain evidence		Human blood/other peripheral tissue evidence		CFG score	GAIN-BP P-value
		WTCC [2007] P-value	NIMH [Baum et al., 2008], P-value	German [Baum et al., 2008], P-value	step-BD [Sklar et al., 2008], P-value	Mouse genetic evidence (QTL, TG)	Mouse models brain evidence [Ogden et al., 2004; Le-Niculescu et al., 2008c] ST AMY (I) ST PFC (D)	Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/other peripheral tissue evidence	Human genetic evidence (linkage or association)	Human postmortem brain evidence	Human blood/other peripheral tissue evidence						
6546	SLC8A1 [Solute carrier family 8 (sodium/calcium exchanger), member 1]	4.57E-03, rs10490049	2.77E-04, rs17025372	2.28E-02, rs381797	7.44E-03, rs12052585	Abnormal emotion/affect behavior (QTL)	Le-Niculescu et al., 2008b	2p22.1										6.5	1.30E-02
8224	SYN3 (Synapsin III)	1.67E-04, rs11089599	4.94E-03, rs130301	4.17E-03, rs3788467	2.03E-02, rs933255													6.5	7.97E-03
7074	TIAM1 (T-cell lymphoma invasion and metastasis 1)	7.39E-05, rs13340018	1.82E-03, rs12482796	2.65E-03, rs2257062	2.49E-03, rs845945	Abnormal emotion/affect behavior (QTL)												6.5	3.31E-02
128553	TSHZ2 (Teashirt family zinc finger 2)	1.98E-02, rs7263115	8.22E-03, rs2741356	3.58E-04, rs169270	1.73E-02, rs6068531	Abnormal emotion/affect behavior (QTL)												6.5	8.16E-03
79683	ZDHHC14 [Zinc finger, DHHC domain containing 14]	4.09E-03, rs1885452	4.59E-03, rs596183	3.56E-02, rs16900254	4.89E-03, rs17297221													6.5	2.40E-02

Top genes with a CFG score of 6.5 and above (n = 56) are shown. The complete list of genes (n = 1,528) is available as Supplementary Information online. I, increased; D, decreased; D, decreased in expression. For human blood data: I, increased in high mood (mania); D, decreased in high mood (mania)/increased in low mood (depression). [For human blood data, where references other than Le-Niculescu et al. [2009a] are cited, the studies are in lymphoblastoid cell lines without correlation with mood state. I, increased; D, decreased]. METH, methamphetamine; VPA, valproate; PFC, prefrontal cortex; AMY, amygdala; CP, caudate putamen; NAC, nucleus accumbens; VT, ventral tegmentum; DBP, knock-out mice; NST, nonstressed; ST, stressed; BP, bipolar disorder; BAD, bipolar affective disorder; MDD, major depressive disorder. TG, transgenic; QTL, quantitative trait locus. For additional human genetic evidence, assoc, association evidence, where that is not mentioned, the evidence is linkage only. Gene symbols underlined are blood biomarker candidate genes. P-values in bold are <0.001. The column on the right of the bolded line depicts replication of findings in an independent bipolar GWAS [GAIN-BP]. Forty-six of our top 56 genes had a P < 0.05 in the GAIN-BP study. The P-value cited is the lowest for any SNP designated by SNPPER to be within the gene or flanking regions. As there were 6,041 genes at P < 0.05 in that study, and the number of genes in the human genome is estimated at 20,500 [Clamp et al., 2007], the enrichment factor provided by our approach is (46/56)/(6,041/20,500) = 2.8-fold. [As a caveat, the GAIN-BP P-values are calculated for the whole cohort in the study, which contains some overlap with the cohort in the NIMH study (see Materials and Methods Section)]. In other words, the positive predictive value (PPV) of a CF6FG score greater than 6 (i.e., > 50% maximum CF6 score) for predicting possible involvement in bipolar disorder for a gene (P < 0.05) is 46/56 = 82.1%. The negative predictive value (NPV) is (20500-56) = 70.7%. Such numbers compare favorably with detection modalities used in science in general, and medicine in particular.

TABLE II. Biological Pathways

Top networks		Score
Associated network functions		
(1) Nervous system development and function, neurological disease, genetic disorder		38
(2) Cellular compromise, neurological disease, drug metabolism		38
(3) Cellular assembly and organization, cardiovascular system development and function, cellular growth and proliferation		24
(4) Amino acid metabolism, cancer, cell morphology		19
Top bio functions		P-Value
Diseases and disorders		# Molecules
Genetic disorder	1.52E-19 to 3.47E-03	53
Neurological disease	1.52E-19 to 3.67E-03	47
Psychological disorders	1.52E-19 to 1.95E-03	35
Endocrine system disorders	4.63E-15 to 3.67E-03	38
Metabolic disease	2.44E-14 to 1.05E-07	38
Molecular and cellular functions		P-Value
Cellular assembly and organization		# Molecules
Cell-to-cell signaling and interaction	4.21E-08 to 3.67E-03	23
Cellular movement	8.46E-08 to 3.67E-03	26
Amino acid metabolism	2.87E-07 to 3.67E-03	15
Molecular transport	1.68E-06 to 2.81E-03	15
Physiological system development and function		P-Value
Behavior		# Molecules
Nervous system development and function	4.22E-13 to 2.94E-03	19
Organismal functions	2.83E-11 to 3.67E-03	33
Tissue morphology	1.97E-10 to 1.45E-09	10
Hematological system development and function	2.27E-05 to 3.67E-03	17
Top canonical pathways		P-Value
		Ratio
(1) G-protein coupled receptor signaling	8.50E-07	8/218 [0.037]
(2) CREB signaling in neurons	3.00E-06	7/194 [0.036]
(3) Synaptic long term depression	1.46E-05	6/164 [0.037]
(4) cAMP-mediated signaling	2.47E-05	6/164 [0.037]
(5) Neuropathic pain signaling in dorsal horn neurons	3.54E-05	5/104 [0.048]

Ingenuity pathway analysis of the top candidate genes from Table III.

genes that had nominal P values <0.05 in one or several of the four GWAS datasets (Wellcome, German, NIMH, STEP-BD) we used were identified. The best P values SNPs in each study were assembled in a GRP panel (Table III), and tested in the GAIN-BP data. As a caveat, not all the SNPs in our GRP panel had been genotyped in the GAIN-BP. Overall, out of 216 SNPs in our panel ($4 \text{ SNPs} \times 56 \text{ genes} = 224 \text{ SNPs}$ theoretically, but some genes did not have a nominally significant SNPs in one or another of the 4 discovery GWAS), only 118 were tested in the GAIN-BP sample.

Each SNP has two alleles (represented by base letters at that position). One of them is associated with the illness (affected), the other not (non-affected), based on the odds ratio from the GAIN-BP. We assigned the affected allele a score of 1 and the non-affected

allele a score of 0. A two-dimensional matrix of subjects by GRP panel alleles is generated, with the cells populated by 0 or 1 (Fig. S3). A SNP in a particular individual subject can have any permutation of 1 and 0 (1 and 1, 0 and 1, 1 and 0, 0 and 0). By adding these numbers, the minimum score for a SNP in an individual subject is 0, and the maximum score is 2. By adding the scores for all the alleles in the panel, averaging that, and multiplying by 100, we generate for each subject an average score corresponding to a genetic loading for disease, which we call Genetic Risk Predictive Score (GRPS). From lower to higher genetic risk, the GRPS has a minimum value of 0 and maximum value of 100. As a caveat, the assignments of 0 and 1 were made based on information for that allele in GAIN-BP for EA subjects, and separately for AA subjects, and rests on the

TABLE III. Intra-Pathway Epistasis (INPEP) Testing Identifies Genes That May Work Together

Ingenuity top canonical pathways	Genes								Nominal epistatic <i>P</i> values
G-Protein coupled receptor signaling	CAMK2A	GRM3	PDE10A*	OPRM1	GNAI1	PRKCE	PDE4B*	HTR2A	*0.0133
CREB signaling in neurons	CAMK2A	GRM3	GRIA1	CREBBP*	GNAI1*	PRKCE	GRIK1		*0.0138
Synaptic long term depression	NOS1*	GRM3*	GRIA1	RYR3	GNAI1	PRKCE			*0.0173
cAMP-mediated signaling	CAMK2A	GRM3	PDE10A*	OPRM1	GNAI1	PDE4B*			*0.0133
Neuropathic pain signaling in dorsal horn neurons	CAMK2A	BDNF	GRM3*	GRIA1	PRKCE*				*0.0298

Inside each of the top canonical pathways depicted in Table II, we tested for epistatic interactions between genes in the pathway, in an independent dataset, the GAIN-BP, as a way of identifying and prioritizing interactions. The top epistatic interactions in each pathway are depicted in bold. These genes merit future follow-up work to elucidate the biological and pathophysiological relevance of their interactions. As a caveat, the *P*-value was not corrected for multiple comparisons.

assumption that the same alleles are associated with bipolar disorder in all subjects of the same ethnicity. However, the GAIN-BP test GWAS is not used to select the panel of genes and SNPs in the GRP panel, which is being derived completely independently from the CFG analysis of the four discovery GWAS.

The software package PLINK (<http://pngu.mgh.harvard.edu/~purcell>) was used to extract individual genotype information for each subject from the GAIN-BP GWAS data files. We analyzed separately European-American (EA) and African-American (AA) bipolar subjects and controls, to examine any potential ethnicity variability (Fig. 3a,b). To test for significance between bipolar and control subjects, a one-tailed *t*-test was performed between the bipolar subjects and the control subjects. We also analyzed males and females separately from each other, to look at any gender-induced variability (Fig. 3c,d). Finally, we tested for the ability of the GRPS to distinguish between bipolar subjects based on an important clinical variable, episode frequency, which is the sum of all episodes of illness (depression and mania), divided by the number of years of illness. We used a case-case design with extremes in phenotype. Thus, compared the GRPS in subjects with the top 1/3 of episode frequency scores versus subjects with the bottom 1/3 of episode frequency scores (Fig. 3e,f).

GRPS Prediction Testing

In a subsequent analysis, we used a split cohort design. We split the GAIN-BP samples for each ethnicity into a 2/3 cohort used for setting GRPS thresholds for bipolar and controls, and a 1/3 cohort used for testing the predictive value of these settings. Inside each ethnicity, the assignment to cohorts was matched for gender, but otherwise random (Table SI).

The average GRPS score for bipolar subjects in the 2/3 cohort is used as a cut-off for bipolar in the test 1/3 cohort (i.e., being above that threshold), and the average GRPS score for controls in the 2/3 cohort is used as a cut-off for controls in the test 1/3 cohort (i.e., being below that threshold). The subjects who are in between these two thresholds are called undetermined. Furthermore, to stratify risk, we categorized subjects in the 1/3 testing cohort into Category 1 if they fall within one standard deviation above the bipolar threshold, and Category 1 if they fall within one standard deviation below the control threshold. Category 2 are between one and two standard

deviations from the thresholds, Category 3 between two and three standard deviations, and Category 4 are those who fall beyond three standard deviations of the threshold. The positive predictive value (PPV) of the test was calculated for each of the categories (Fig. 4).

RESULTS

Top Candidate Genes


In order to minimize false negatives, we initially cast a wide net, using as a filter a minimal requirement for a gene to have both some genetic and some functional genomic evidence. We thus generated an initial list of 1,657 unique genes with at least a SNP at $P < 0.05$ in at least one of the four primary GWAS analyzed, that also had some functional (gene expression) evidence (human or animal model data), implicating them in bipolar disorder or depression. Of interest, a previous similar analysis by us using just three GWAS [Le-Niculescu et al., 2009b] yielded 1,529 unique genes, suggesting that: (1) with our genetic-genomic filtering of the GWAS in the primary analysis we are already capturing most of the genes that may be involved in bipolar disorder, with additional studies providing an asymptotic contribution beyond this point; and (2) that, using our thresholds and minimal requirements, the number of genes potentially involved, directly or indirectly, in bipolar disorder may be indeed quite large, up to 10% of the genome (see also Supplementary Information—Fig. S1).

In order to minimize false positives, we then used a CFG analysis integrating multiple lines of evidence to prioritize this initial list of 1,657 genes, and focused our subsequent analyses on only the top CFG scoring candidate genes. 56 genes had a CFG score of above 6 (>50% of maximum possible score) (Table III and Fig. 2).

As a way of testing the validity of our approach, we have examined whether our top findings were over-represented in an independent GWAS of bipolar disorder, the GAIN-BP study. Forty-six of the top 56 genes identified by our approach had a *P*-value of < 0.05 in that independent study, an estimated almost threefold enrichment over what would be expected by chance alone in that study (see Table III).

Candidate Blood Biomarkers

Of the top candidate genes from Table III (see also Fig. 2), 22 out of 56 have prior blood gene expression evidence implicating them as



potential blood biomarkers. The additional evidence provided by GWAS data suggests a genetic rather than purely environmental (medications, stress) basis for their alteration in disease, and their potential utility as trait rather than purely state markers.

Biological Pathways

Ingenuity pathway analysis was carried out on the top 56 genes, revealing results similar to our previous work [Le-Niculescu et al., 2009b]. Notably, G-protein coupled receptor signaling, cAMP related signaling and synaptic long-term depression were the top canonical pathways over-represented in bipolar disorder, which is informative (and reassuring, as these pathways are highly druggable) for new drug discovery efforts by pharmaceutical companies.

Intra-Pathway Testing for Epistasis (INPEP)

Epistatic interactions testing inside each of these pathways, using the independent GAIN-BP data, revealed some nominally significant pair-wise *P* values (Table II). For example, the possible interactions between CREBBP and GNAI1, and between NOS1 and GRM3 are non-obvious.

These prioritized pairs of genes inside each pathway may merit future hypothesis driven confirmatory genetic studies in independent cohorts, as well as testing for mechanistic interactions relevant to bipolar disorder pathophysiology in follow-up biological studies, such as transgenic mice studies.

DISCUSSION

Our CFG approach helped prioritize, as in our previous work [Le-Niculescu et al., 2009b] and as expected, genes for which there was consistent evidence among the four discovery GWAS datasets, or

stronger evidence in one or another of the datasets. However, it also prioritized genes with weaker evidence in the GWAS data, but with strong independent evidence in terms of gene expression studies and other prior human or animal genetic work.

At the very top of our list of candidate genes for bipolar disorder we have five genes: ARNTL, MBP, BDNF, NRG1, and RORB.

Aryl hydrocarbon receptor nuclear translocator-like (ARNTL), a transcription factor, is a circadian clock gene. Another circadian top candidate genes identified by our analysis is RORB (Fig. 2 and Table III). RORB was also recently reported by us to be associated with bipolar disorder in an independent pediatric bipolar sample [McGrath et al., 2009]. Circadian rhythm and sleep abnormalities have long been described in bipolar disorder—excessive sleep in the depressive phase, reduced need for sleep in the manic phase [Bauer et al., 2006]. Sleep deprivation is one of the more powerful and rapid acting treatment modalities for severe depression, and can lead to precipitation of manic episodes in bipolar patients [Wirz-Justice et al., 2004]. We had previously described the identification of clock gene D-box binding protein (DBP) as a potential candidate gene for bipolar disorder [Niculescu et al., 2000b], using a CFG approach. DBP was changed in expression by acute methamphetamine treatment in rat pre-frontal cortex (PFC) [Niculescu et al., 2000b], and mapped near a human genetic linkage locus for bipolar disorder [Morissette et al., 1999] and for depression [Zubenko et al., 2002] on chromosome 19q13. Subsequently, DBP was also reported changed in expression by acute and chronic amphetamine treatments in mice [Sokolov et al., 2003]. Moreover, DBP knock-out mice have abnormal circadian and homeostatic aspects of sleep regulation [Franken et al., 2000]. More recently, we have conducted extensive behavioral and gene expression studies in DBP KO mice. These mice display a bipolar-like phenotype [Le-Niculescu et al., 2008c], which is modulated by stress. Decreases in DBP expression have also been recently reported in fibroblasts from bipolar subjects [Yang et al., 2008]. In parallel, work carried out by us using an expanded CFG approach in a mouse pharmacogenomic model for bipolar disorder identified ARNTL and a series of other clock genes (CRY2, CSNK1Ds, and CCR4/nocturnin), as potential bipolar candidate genes [Ogden et al., 2004]. Following that, three independent reports have shown some suggestive association for ARNTL in human bipolar samples [Mansour et al., 2006; Nievergelt et al., 2006; Shi et al., 2008]. ARNTL is upstream of DBP in the circadian clock intracellular molecular machinery, driving the transcription of DBP [Ripperger and Schibler, 2006; van der Veen et al., 2006]. An increase in ARNTL gene expression was reported in postmortem brains from bipolar subjects [Nakatani et al., 2006]. Seasonal affective disorder (SAD), a variant of bipolar disorder [Magnusson and Partonen, 2005], is tied to the amount of daylight, which is a primary regulator of circadian rhythms and clock gene expression; associations between polymorphisms in the clock genes ARNTL, PER2, and NPAS2 and SAD have previously been reported [Johansson et al., 2003; Partonen et al., 2007]. Overall, ARNTL, RORB, DBP and related circadian clock genes are compelling candidates for involvement in bipolar disorders, acting as rheostats as well as underlying the core clinical phenomenology of cycling and switching from depression to mania [Bunney and Bunney, 2000; Wager-Smith and Kay, 2000; Niculescu et al., 2000b; Niculescu and Kelsoe, 2001; Kelsoe and Niculescu, 2002; Lenox et al., 2002; Hasler

et al., 2006; Wirz-Justice, 2006; McClung, 2007; Le-Niculescu et al., 2008c].

Myelin basic protein (MBP) is involved in white matter build-up and connectivity processes [Harauz et al., 2009]. Based on human postmortem and blood biomarker work, as well as animal models (see Table III), it may be decreased in expression in the depressive phase of bipolar disorder, leading to a slowing of action potential transmission, potential disconnection between brain regions, and outward psychomotor retardation. The additional evidence provided by GWAS data indicates a genetic rather than purely environmental (medications, stress) basis for its alteration in disease, and its potential utility as trait marker for increased vulnerability. MBP alterations have also been reported in other neuropsychiatric disorders such as schizophrenia [Parlapani et al., 2009], alcoholism [Lewohl et al., 2005], multiple sclerosis [Zamvil et al., 1985], as well as an animal models of stress reactivity [Le-Niculescu et al., 2008c]. Myelin-related genes may be a common if non-specific denominator of vulnerability to mental illness in response to stress [Le-Niculescu et al., 2008c].

BDNF is a growth factor involved in neurotrophicity and synaptic transmission. Other growth factor top candidate genes identified by our analysis include NRG1 and PTN (Fig. 2 and Table III). BDNF has been previously implicated in a variety of neuropsychiatric disorders, by both animal model and human studies: depression [Pezawas et al., 2008; Sen et al., 2008], bipolar disorder [Ogden et al., 2004], anxiety, alcoholism [Rodd et al., 2007], and schizophrenia [Le-Niculescu et al., 2007a; Chao et al., 2008]. Notably, there are several candidate gene association studies to date implicating BDNF in bipolar disorder [Fan and Sklar, 2008; Liu et al., 2008].

Amyloid beta precursor protein (APP), an Alzheimer disease (AD) candidate gene, is among the top candidate gene for bipolar disorder (Table III). Another key gene involved in AD, GSK3b, is also present on our list of top candidate genes. Previous epidemiological literature has pointed to increased AD in bipolar patients, and the prophylactic effect of the mood stabilizer lithium on the incidence of AD in bipolar patients [Nunes et al., 2007]. Notably, GSK3b is a target of lithium treatment [Beaulieu et al., 2008a], as well as of serotonergic anti-depressants [Beaulieu et al., 2008b]. APP has recently been shown to have a neurotrophic role [Oh et al., 2008], similar to growth factors such as BDNF. APP has also been reported to be increased in expression in bipolar postmortem brains compared to normal controls [Jurata et al., 2004]. It remains to be seen if APP's role in AD is pathogenic or is in fact a defense/compensatory mechanism to try to maintain neuronal survival [Rohn et al., 2008]. The possibility that drugs that regulate APP levels may have an impact on mood (i.e., downregulation of APP may be depressogenic) needs to be explored, given the prevalence of depression in the elderly in general [Alexopoulos et al., 2005], and in AD patients in particular [Sun et al., 2008]. In any case, this is an intriguing example of potential genetic co-morbidity, overlap and interdependence between mood and cognition.

Genetic Risk Prediction

Once the genes involved in a disorder are identified, and prioritized for likelihood of involvement, then an obvious next step is

developing a way of applying that knowledge to genetic testing of individuals to determine risk for the disorder. Based on our comprehensive identification of top candidate genes described above, we pursued a polygenic panel approach, with digitized binary scoring for presence or absence, similar to the one we have devised and employed in the past for biomarkers [Le-Niculescu et al., 2009a]. Somewhat similar approaches, looking however at larger panels of markers without CFG prioritization, were subsequently also described by other groups [Purcell et al., 2009].

We have chosen the best SNPs in our CFG prioritized genes by their *P* values in the GWAS datasets used, and assembled a genetic risk prediction (GRP) panel out of those SNPs (Table III). We then developed a genetic risk prediction score (GRPS) for bipolar disorder based on the presence or absence of the alleles of the SNPs associated with the illness, and tested the GRPS in an independent study (GAIN-BP) for which we had both genotypic and clinical data available, comparing the bipolar subjects to demographically matched normal controls (Fig. 3).

We demonstrate that in independent test cohorts, the GRPS differentiates between subjects with bipolar disorder and normal controls, in both European-American (EA) and African-American (AA) subjects (Fig. 3a,b). The GRPS also differentiates between high episode frequency and low episode frequency bipolar subjects in EA, but not AA subjects (Fig. 3e,f). Gender analyses exhibited slight trends towards higher GRPS in males than females in both ethnicities, that did not reach statistical significance (Fig. 3c,d). Lastly, we also describe a prototype of how such testing could be used at an individual rather than population level, to categorize individuals by risk and aid diagnostic and personalized medicine approaches (Fig. 4).

Our results show that a relatively small size panel identified by CFG analysis can differentiate very well between bipolar disorder subjects and controls at a population level, although at an individual level the margin is razor thin (Figs. 3 and 4). On average, a bipolar subject differs from a control subject by about 2 alleles out of 236 tested. The latter point suggests that the cumulative combinatorics of common gene variants plays a major role in genetic risk for illness. Overall, our work sheds light on the genetic architecture and pathophysiology of bipolar disorder. In particular, it has implications for genetic testing to assess risk for illness. Our evaluation of the predictive value of the GRPS suggests some utility by itself at identifying risk for illness (Fig. 4). More likely, such genetic information will have to be combined with family history and other clinical information (phenomics) [Niculescu et al., 2006], as well as with blood biomarker testing [Le-Niculescu et al., 2009a], to provide a comprehensive picture of risk of illness [Niculescu, 2006; Niculescu et al., 2009].

Limitations and Confounds

No correction of best *P* values for number of SNPs tested/gene size effect was performed. While this is arguably a valid statistical issue for genetic studies by themselves, some of the multiple SNPs tested per gene could be in linkage disequilibrium, and the Bonferroni correction might be too conservative [Rice et al., 2008]. One would expect some noise due to gene size, as larger genes have more SNPs tested per gene. However, we did not observe a significant correlation between gene size and our top candidate gene prioritization using CFG (Supplementary Information—Fig. S2). That may be




FIG. 3. The genetic risk prediction score (GRPS) for bipolar disorder differentiates between bipolar subjects and normal controls in an independent study, in two different ethnic groups. The GRPS is based on a panel of the best *P*-value SNPs (*n* = 216) from the best top genes (*n* = 56) for bipolar disorder identified by CFG of four GWAS for bipolar disorder (see Table III). The GRPS shows statistically significant differences between subjects with bipolar disorder and normal controls, in European American (EA) [a] as well as in African Americans (AA) [b], in an independent GWAS study [GAIN-BP, Smith et al., 2009]. Out of 216 SNPs in our panel, 118 SNPs were genotyped in the GAIN-BP study. Gender analyses exhibited slight trends towards higher GRPS in males than females in both ethnicities, that did not reach statistical significance [c,d]. Episode frequency: of note, the GRPS is able to differentiate between high episode frequency and low episode frequency bipolar subjects in EA [e], but not AA subjects [f]. Episode frequency is a measure of clinical severity, tied to recurrence and to cycling between euthymia, depression and mania.

due to the fact that we are using this evidence for integration across platforms and modalities, along with a series of other lines of evidence that have their own attendant noise, as part of a Bayesian-like approach to pull signal from noise and prioritize findings. The

convergence of lines of evidence arguably factors out the noise of the different individual approaches, and makes our network-like CFG approach relatively resilient to error even when one or another of the nodes (lines of evidence) is weak (Fig. 1).




FIG. 4. Prototype of how GRPS testing could be used at an individual rather than population level, to aid diagnostic and personalized medicine approaches. We split the GAIN-BP samples from each ethnicity into a 2/3 cohort used for setting GRPS thresholds for bipolar and controls (a,c), and a 1/3 cohort used for testing the predictive value of these settings (b,d). The average GRPS score for bipolar subjects in the 2/3 cohort is used as a cut-off for bipolar in the test 1/3 cohort (i.e., being above that threshold), and the average GRPS score for controls in the 2/3 cohort is used as a cut-off for controls in the test 1/3 cohort (i.e., being below that threshold). The subjects who are in between these two thresholds are called undetermined. Furthermore, to stratify risk, we categorized subjects in the 1/3 testing cohort into Category 1 if they fall within one standard deviation above the bipolar threshold, and Category 1 if they fall within one standard deviation below the control threshold. Category 2 subjects are between one and two standard deviations from the thresholds, Category 3 between two and three standard deviations, and Category 4 are those who fall beyond three standard deviations of the threshold. The positive predictive value (PPV) of the tests increases in the higher categories, and the test is somewhat better at distinguishing controls [i.e., in a practical application, individuals that are lower risk of developing the illness] than bipolars [i.e., in a practical application, individuals that are higher risk of developing the illness].

Our approach relies on a list of genes from the GWAS datasets generated by SNPPER identifying SNPs in genes. We may thus be missing genes where the assignment is not made by the software, and discarding SNPs that fall into intergenic or regulatory regions, such as promoter or enhancer regions. Moreover, genes where the illness associated SNPs do not lead to a change in expression levels are not included in our CFG-GWA cross-validation. Similarly, genes that have changes in expression levels but no intragenic SNP

in the GWAS datasets are not included. Interestingly, some of these latter genes may be changed in expression as a consequence of distal regulatory SNPs or other genes in a network, an exciting area for future systems biology studies awaiting better bioinformatic tools and data analysis now on the horizon [Stumpf et al., 2008]. Our panel of genes prioritized by CFG is certainly not exhaustive, it is just an example of one approach. Some of the genes with strong published evidence of association, such as ANK3 [Ferreira et al.,

2008a; Schulze et al., 2009] and CACNA1C [Ferreira et al., 2008a], are not prioritized by our approach, (although related genes, ANK2 and CACNA1A, are) (Table III). It may be that genes prioritized by *P* values in genetic studies alone are the result of a *fit-to-cohort* phenomenon, resulting in poor reproducibility and predictive value in independent cohorts [Niculescu and Le-Niculescu, 2010; Paynter et al., 2010]. The few of them that are reproduced across studies may be more of a common denominator in bipolar patients (which tend to be heterogeneous), somewhat like house-keeping genes, but not necessarily more biologically relevant or important. CFG arguably identifies and prioritizes genes that have functional evidence and hence are more likely to be biologically relevant [Niculescu and Le-Niculescu, 2010]. By being in essence a *fit-to-disease* approach, CFG also generates findings that are more reproducible and have predictive value in independent studies and cohorts, as we have demonstrated in our previous work on blood biomarkers [Kurian et al., 2009; Le-Niculescu et al., 2009a], and as we demonstrate in this current genetic work. That is the key litmus test, in our view.

Other animal models data could potentially be used for CFG cross-validation, in addition to the data from the pharmacogenomic (methamphetamine/valproate) [Ogden et al., 2004] and the genetic (DBP knock-out mouse) [Le-Niculescu et al., 2008c] models that we generated and used. However, these are some of the best animal models with corresponding comprehensive brain and blood gene expression datasets published to date. Moreover, we relied, as an additional line of evidence, on an extensive public mouse QTL/transgenic database.

As new human blood, postmortem brain, and human genetic studies are published, new evidence will be available for some of the genes we have identified. However, any new evidence will likely not remove genes from our results, but rather move them up higher in the prioritization list/pyramid (Fig. 2).

Different ways of weighing the lines of evidence included in the CFG analysis rather than the equal weight approach we have used may become available in the future, based on more empirical and quantitative methods. Other ways of weighing the scores of line of evidence may give slightly different results in terms of prioritization, if not in terms of the list of top genes per se.

Pathways identified by Ingenuity may be based on some of the same body of knowledge and published literature used in our direct CFG scoring. However, it is reassuring to see that different independent systematization and curation efforts lead to a consistent picture of genes involved in behavior, neurological disease, psychological disorders, and nervous system development coming up at the top of the over-represented pathways from our top candidate genes for bipolar disorder identified by our genetic-genomic combined approach.

CONCLUSIONS AND FUTURE DIRECTIONS

First, in spite of these limitations, our analysis is arguably the most comprehensive integration of genetics and functional genomics to date in the field of bipolar disorder, yielding a series of candidate genes, blood biomarkers, pathways and mechanisms, that are prime targets for follow-up hypothesis driven studies. Such studies may

include individual candidate gene association studies with more SNPs tested per gene, deep re-sequencing, and/or biological validation such as cell culture [Pletnikov et al., 2007] and transgenic animal work [Hikida et al., 2007; Le-Niculescu et al., 2008c].

Second, our work provides additional integrated evidence focusing attention and prioritizing a number of genes as candidate blood biomarkers for bipolar disorder, with an inherited genetic basis (Table III). While prior evidence existed as to alterations in gene expression levels of those genes in whole-blood samples or lymphoblastoid cell lines (LCLs) from mood disorders patients, it was unclear prior to our analysis whether those alterations were truly related to the disorder or were instead related to medication effects and environmental factors.

Third, our work provides proof how a combined approach, integrating functional and genotypic data, can be used for other complex disorders-psychiatric and non-psychiatric. What we are seeing across GWAS of complex disorders are not necessarily the same genes showing the strongest signal, but rather consistency at the level of gene families or biological pathways. The distance from genotype to phenotype may be a bridge too far for genetic-only approaches, given the intervening complex layers of epigenetics, gene expression regulation and endophenotypes [Tan et al., 2008]. Using GWAS data in conjunction with gene expression data as part of CFG or integrative genomics [Degnan et al., 2008] approaches, followed by pathway-level analysis of the prioritized candidate genes, can lead to the unraveling of the genetic code of complex disorders such as bipolar disorder.

Fourth, we have focused attention on key biological pathways in bipolar disorder, and used genetic epistatic testing to identify and prioritize molecular interactions inside those pathways. We believe that this intra-pathway epistasis testing approach (INPEP) may help with future work aimed at dissecting the molecular architecture of complex disorders.

Fifth, we have put together a panel of best *P* value single nucleotide polymorphisms (SNPs), based on the top candidate genes we identified. Such a panel could be used for genetic testing for bipolar disorder. To that end, we have developed a genetic risk prediction score (GRPS) based on our panel, and demonstrate how in independent cohorts, the GRPS differentiates between patients with bipolar disorder and normal controls. Based on the GRPS, we demonstrate a prototype of individual subject categorization for risk of illness. We anticipate that the GRPS approach will have utility for other complex disorders, psychiatric and non-psychiatric.

Lastly, while we cannot exclude that rare genetic variants with major effects may exist in some individuals and families, we propose a *cumulative combinatorics of common variants* genetic model for bipolar disorder based on our findings (Fig. 5), to account for the razor thin genetic load margin between clinically ill subjects and normal controls, which leaves a major role to be played by the environment [Lahiri et al., 2009; Niculescu et al., 2009]. A stressful/hostile environment may lead to sub-threshold illness even in normal genetic load individuals, whereas a favorable environment may lead to supra normative functioning in certain life areas for individuals who carry a higher genetic risk. From a speculative standpoint, this proposed flexible interplay between genetic load, environment and phenotype may permit evolution to engender




FIG. 5. The genetic architecture of bipolar disorders: Cumulative combinatorics of common gene variants and environment [CC × CGV × E] model. We proposed in our previous work [Le-Niculescu et al., 2009b] that the repertoire of genes that may be involved directly or indirectly in bipolar disorders/mood regulation is large, up to 10% of the genes in the genome (complexity). Our current work suggests that different combinations of genes/alleles are found in different individuals (heterogeneity), and many alleles in these genes are shared between bipolars and controls (overlap). The environment may play a key interactive role in the trajectory from genetic risk to ultimate phenotype (interdependence), by modulating gene expression. For example, with the right environment a higher genetic load (GRPS score) individual may become normal or even a high performer. This shuffling of the genetic deck of cards and the interaction with the environment provide a basis for Darwinian adaptation and evolution of mood, a key bodily function synchronizing energy metabolism, trophicity and activity to external and internal milieu conditions [Le-Niculescu et al., 2009a,b]. Circadian clock molecular mechanisms, involving ARNTL, RORB, DBP and other molecules, may be essential mediators [Niculescu et al., 2000b; Ogden et al., 2004; Takahashi et al., 2008; Le-Niculescu et al., 2008c; Zhang et al., 2009]. Geometric symbols in the figure depict different genes/alleles.

diversity, select and conserve alleles, ultimately shaping population groups.

From a pragmatic utility standpoint, we would like to suggest that genetic testing with highly prioritized panels of best markers will have, by itself, a rather modest role in informing clinical decisions regarding early intervention and prevention efforts, for example before the illness fully manifests itself clinically, in young offspring from high-risk families. After the illness manifests itself, biomarker and phenomic testing approaches, including clinical data, may have higher yield than genetic testing, and a multi-modal integration of testing modalities may be optimal, as individual markers are likely to not be specific for a single disorder. The continuing re-evaluation in psychiatric nosology [Niculescu et al., 2009; O'Donovan et al., 2009] brought about by recent advances

will have to be taken into account as well for final interpretation of any such testing. Our emerging appreciation of the complexity, heterogeneity, overlap and interdependence of major psychiatric disorders as currently defined, and their building blocks/Lego-like nature [Le-Niculescu et al., 2007a], may make the development of tests for specific modular disease manifestations (mood, psychosis, anxiety) [Niculescu et al., 2009] more useful and precise than those for broad diagnostic categories like bipolar disorder, schizophrenia or post-traumatic stress disorder.

ACKNOWLEDGMENTS

This work was supported by funds from INGEN (Indiana Genomics Initiative of Indiana University), INBRAIN (Indiana Center for Biomarker Research In Neuropsychiatry), NARSAD Young Investigator Award and VA Merit Award to ABN, as well as NIMH R01 MH071912-01 to Ming Tsuang and ABN. ABN would like to thank Nicholas Schork for extensive discussions on genetic data analyses, Daniel Salomon, Sunil Kurian and Howard Edenberg for help and advice with microarray data analyses, Ming Tsuang and Steven Faraone for help and advice with translational studies, as well as Mariano Erpe, Joyti Gupta and Jesse Townes for their precise work with database maintenance and data analyses. Most importantly, we would like to thank our coworkers in the field whose painstaking work we have cited and integrated in our analyses, particularly the BiGS consortium (see Supplementary Information), as well as the subjects who participated in these studies, their families and their caregivers. Without their contribution, such work to advance the understanding of mental illness would not be possible. This work is, in essence, a field-wide collaboration.

REFERENCES

- Aguilera M, Arias B, Wichers M, Barrantes-Vidal N, Moya J, Villa H, van Os J, Ibanez MI, Ruiperez MA, Ortet G, et al. 2009. Early adversity and 5-HTT/BDNF genes: New evidence of gene-environment interactions on depressive symptoms in a general population. *Psychol Med* 39(9): 1425–1432.
- Alexopoulos GS, Schultz SK, Lebowitz BD. 2005. Late-life depression: A model for medical classification. *Biol Psychiatry* 58(4):283–289.
- Amano K, Yamada K, Iwayama Y, Detera-Wadleigh SD, Hattori E, Toyota T, Tokunaga K, Yoshikawa T, Yamakawa K. 2008. Association study between the Down syndrome cell adhesion molecule (DSCAM) gene and bipolar disorder. *Psychiatr Genet* 18(1):1–10.
- Arai M, Itokawa M, Yamada K, Toyota T, Arai M, Haga S, Ujike H, Sora I, Ikeda K, Yoshikawa T. 2004. Association of neural cell adhesion molecule 1 gene polymorphisms with bipolar affective disorder in Japanese individuals. *Biol Psychiatry* 55(8):804–810.
- Arranz MJ, Erdmann J, Kirov G, Rietschel M, Sodhi M, Albus M, Ball D, Maier W, Davies N, Franzek E, et al. 1997. 5-HT2A receptor and bipolar affective disorder: Association studies in affected patients. *Neurosci Lett* 224(2):95–98.
- Aston C, Jiang L, Sokolov BP. 2005. Transcriptional profiling reveals evidence for signaling and oligodendroglial abnormalities in the temporal cortex from patients with major depressive disorder. *Mol Psychiatry* 10(3):309–322.
- Atz ME, Rollins B, Vawter MP. 2007. NCAM1 association study of bipolar disorder and schizophrenia: Polymorphisms and alternatively spliced

- isoforms lead to similarities and differences. *Psychiatr Genet* 17(2): 55–67.
- Badenhop RF, Moses MJ, Scimone A, Mitchell PB, Ewen-White KR, Rosso A, Donald JA, Adams LJ, Schofield PR. 2002. A genome screen of 13 bipolar affective disorder pedigrees provides evidence for susceptibility loci on chromosome 3 as well as chromosomes 9, 13 and 19. *Mol Psychiatry* 7(8):851–859.
- Bailer U, Leisch F, Meszaros K, Lenzinger E, Willinger U, Strobl R, Heiden A, Gebhardt C, Doge E, Fuchs K, et al. 2002. Genome scan for susceptibility loci for schizophrenia and bipolar disorder. *Biol Psychiatry* 52(1):40–52.
- Bauer M, Grof P, Rasgon N, Bschor T, Glenn T, Whybrow PC. 2006. Temporal relation between sleep and mood in patients with bipolar disorder. *Bipolar Disord* 8(2):160–167.
- Baum AE, Akula N, Cabanero M, Cardona I, Corona W, Klemens B, Schulze TG, Cichon S, Rietschel M, Nothen MM, et al. 2008. A genome-wide association study implicates diacylglycerol kinase eta (DGKH) and several other genes in the etiology of bipolar disorder. *Mol Psychiatry* 13(2):197–207.
- Beaulieu JM, Marion S, Rodriguiz RM, Medvedev IO, Sotnikova TD, Ghisi V, Wetsel WC, Lefkowitz RJ, Gainetdinov RR, Caron MG. 2008a. A beta-arrestin 2 signaling complex mediates lithium action on behavior. *Cell* 132(1):125–136.
- Beaulieu JM, Zhang X, Rodriguiz RM, Sotnikova TD, Cools MJ, Wetsel WC, Gainetdinov RR, Caron MG. 2008b. Role of GSK3 beta in behavioral abnormalities induced by serotonin deficiency. *Proc Natl Acad Sci USA* 105(4):1333–1338.
- Begemann M, Sargin D, Rossner MJ, Bartels C, Theis F, Wichert SP, Stender N, Fischer B, Sperling S, Stawicki S, et al. 2008. Episode-specific differential gene expression of peripheral blood mononuclear cells in rapid cycling supports novel treatment approaches. *Mol Med* 14(9–10): 546–552.
- Benedetti F, Bernasconi A, Lorenzi C, Pontiggia A, Serretti A, Colombo C, Smeraldi E. 2004. A single nucleotide polymorphism in glycogen synthase kinase 3-beta promoter gene influences onset of illness in patients affected by bipolar disorder. *Neurosci Lett* 355(1–2):37–40.
- Benes FM, Matzilevich D, Burke RE, Walsh J. 2006. The expression of proapoptosis genes is increased in bipolar disorder, but not in schizophrenia. *Mol Psychiatry* 11(3):241–251.
- Bertram I, Bernstein HG, Lendeckel U, Bukowska A, Dobrowolny H, Keilhoff G, Kanakis D, Mawrin C, Bielau H, Falkai P, et al. 2007. Immunohistochemical evidence for impaired neuregulin-1 signaling in the prefrontal cortex in schizophrenia and in unipolar depression. *Ann NY Acad Sci* 1096:147–156.
- Bonnier B, Gorwood P, Hamon M, Sarfati Y, Boni C, Hardy-Bayle MC. 2002. Association of 5-HT(2A) receptor gene polymorphism with major affective disorders: The case of a subgroup of bipolar disorder with low suicide risk. *Biol Psychiatry* 51(9):762–765.
- Brezo J, Bureau A, Merette C, Jomphe V, Barker ED, Vitaro F, Hebert M, Carbonneau R, Tremblay RE, Turecki G. 2009. Differences and similarities in the serotonergic diathesis for suicide attempts and mood disorders: A 22-year longitudinal gene-environment study. *Mol Psychiatry*. [Epub ahead of print]
- Bunney WE, Bunney BG. 2000. Molecular clock genes in man and lower animals: Possible implications for circadian abnormalities in depression. *Neuropsychopharmacology* 22(4):335–345.
- Chagnon YC, Merette C, Bouchard RH, Emond C, Roy MA, Maziade M. 2004. A genome wide linkage study of obesity as secondary effect of antipsychotics in multigenerational families of eastern Quebec affected by psychoses. *Mol Psychiatry* 9(12):1067–1074.
- Chambers JS, Perrone-Bizzozero NI. 2004. Altered myelination of the hippocampal formation in subjects with schizophrenia and bipolar disorder. *Neurochem Res* 29(12):2293–2302.
- Chao HM, Kao HT, Porton B. 2008. BDNF Val66Met variant and age of onset in schizophrenia. *Am J Med Genet Part B* 147B(4):505–506.
- Chee IS, Lee SW, Kim JL, Wang SK, Shin YO, Shin SC, Lee YH, Hwang HM, Lim MR. 2001. 5-HT2A receptor gene promoter polymorphism-1438A/G and bipolar disorder. *Psychiatr Genet* 11(3):111–114.
- Cheng R, Juo SH, Loth JE, Nee J, Iossifov I, Blumenthal R, Sharpe L, Kanyas K, Lerer B, Lilliston B, et al. 2006. Genome-wide linkage scan in a large bipolar disorder sample from the National Institute of Mental Health genetics initiative suggests putative loci for bipolar disorder, psychosis, suicide, and panic disorder. *Mol Psychiatry* 11(3):252–260.
- Choudary PV, Molnar M, Evans SJ, Tomita H, Li JZ, Vawter MP, Myers RM, Bunney WE Jr, Akil H, Watson SJ, et al. 2005. Altered cortical glutamatergic and GABAergic signal transmission with glial involvement in depression. *Proc Natl Acad Sci USA* 102(43):15653–15658.
- Cichon S, Schmidt-Wolf G, Schumacher J, Muller DJ, Hurter M, Schulze TG, Albus M, Borrmann-Hassenbach M, Franzek E, Lanczik M, et al. 2001. A possible susceptibility locus for bipolar affective disorder in chromosomal region 10q25–q26. *Mol Psychiatry* 6(3):342–349.
- Clamp M, Fry B, Kamal M, Xie X, Cuff J, Lin MF, Kellis M, Lindblad-Toh K, Lander ES. 2007. Distinguishing protein-coding and noncoding genes in the human genome. *Proc Natl Acad Sci USA* 104(49): 19428–19433.
- Curtis D, Kalsi G, Brynjolfsson J, McInnis M, O'Neill J, Smyth C, Moloney E, Murphy P, McQuillin A, Petursson H, et al. 2003. Genome scan of pedigrees multiply affected with bipolar disorder provides further support for the presence of a susceptibility locus on chromosome 12q23–q24, and suggests the presence of additional loci on 1p and 1q. *Psychiatr Genet* 13(2):77–84.
- Degnan JH, Lasky-Su J, Raby BA, Xu M, Molony C, Schadt EE, Lange C. 2008. Genomics and genome-wide association studies: An integrative approach to expression QTL mapping. *Genomics*
- Detera-Wadleigh SD, Badner JA, Berrettini WH, Yoshikawa T, Goldin LR, Turner G, Rollins DY, Moses T, Sanders AR, Karkera JD, et al. 1999. A high-density genome scan detects evidence for a bipolar-disorder susceptibility locus on 13q32 and other potential loci on 1q32 and 18p11.2. *Proc Natl Acad Sci USA* 96(10):5604–5609.
- Etain B, Mathieu F, Rietschel M, Maier W, Albus M, McKeon P, Roche S, Kealey C, Blackwood D, Muir W, et al. 2006. Genome-wide scan for genes involved in bipolar affective disorder in 70 European families ascertained through a bipolar type I early-onset proband: Supportive evidence for linkage at 3p14. *Mol Psychiatry* 11(7):685–694.
- Ewald H, Flint T, Kruse TA, Mors O. 2002. A genome-wide scan shows significant linkage between bipolar disorder and chromosome 12q24.3 and suggestive linkage to chromosomes 1p22–21, 4p16, 6q14–22, 10q26 and 16p13.3. *Mol Psychiatry* 7(7):734–744.
- Fallin MD, Lasseter VK, Wolyniec PS, McGrath JA, Nestadt G, Valle D, Liang KY, Pulver AE. 2004. Genomewide linkage scan for bipolar-disorder susceptibility loci among Ashkenazi Jewish families. *Am J Hum Genet* 75(2):204–219.
- Fallin MD, Lasseter VK, Avramopoulos D, Nicodemus KK, Wolyniec PS, McGrath JA, Steel G, Nestadt G, Liang KY, Hagan RL, et al. 2005. Bipolar I disorder and schizophrenia: A 440-single-nucleotide polymorphism screen of 64 candidate genes among Ashkenazi Jewish case-parent trios. *Am J Hum Genet* 77(6):918–936.
- Fan J, Sklar P. 2008. Genetics of bipolar disorder: Focus on BDNF Val66Met polymorphism. *Novartis Found Symp* 289:60–72, discussion 72–73, 87–93.

- Fatemi SH, Reutiman TJ, Folsom TD, Lee S. 2008. Phosphodiesterase-4A expression is reduced in cerebella of patients with bipolar disorder. *Psychiatr Genet* 18(6):282–288.
- Ferreira MA, O'Donovan MC, Meng YA, Jones IR, Ruderfer DM, Jones L, Fan J, Kirov G, Perlis RH, Green EK, et al. 2008a. Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. *Nat Genet* 40(9):1056–1058.
- Ferreira MA, O'Donovan MC, Meng YA, Jones IR, Ruderfer DM, Jones L, Fan J, Kirov G, Perlis RH, Green EK, et al. 2008b. Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. *Nat Genet*
- Franken P, Lopez-Molina L, Marcacci L, Schibler U, Tafti M. 2000. The transcription factor DBP affects circadian sleep consolidation and rhythmic EEG activity. *J Neurosci* 20(2):617–625.
- Freimer NB, Reus VI, Escamilla MA, McInnes LA, Spesny M, Leon P, Service SK, Smith LB, Silva S, Rojas E, et al. 1996. Genetic mapping using haplotype, association and linkage methods suggests a locus for severe bipolar disorder (BPI) at 18q22-q23. *Nat Genet* 12(4):436–441.
- Georgieva L, Dimitrova A, Ivanov D, Nikolov I, Williams NM, Grozeva D, Zaharieva I, Toncheva D, Owen MJ, Kirov G, et al. 2008. Support for neuregulin 1 as a susceptibility gene for bipolar disorder and schizophrenia. *Biol Psychiatry*. 64(5):419–427.
- Green EK, Raybould R, Macgregor S, Gordon-Smith K, Heron J, Hyde S, Grozeva D, Hamshere M, Williams N, Owen MJ, et al. 2005. Operation of the schizophrenia susceptibility gene, neuregulin 1, across traditional diagnostic boundaries to increase risk for bipolar disorder. *Arch Gen Psychiatry* 62(6):642–648.
- Harauz G, Ladizhansky V, Boggs JM. 2009. Structural polymorphism and multifunctionality of myelin basic protein. *Biochemistry* 48(34):8094–8104.
- Hasler G, Drevets WC, Gould TD, Gottesman II, Manji HK. 2006. Toward constructing an endophenotype strategy for bipolar disorders. *Biol Psychiatry*. 60(2):93–105.
- Hikida T, Jaaro-Peled H, Seshadri S, Oishi K, Hookway C, Kong S, Wu D, Xue R, Andrade M, Tankou S, et al. 2007. Dominant-negative DISC1 transgenic mice display schizophrenia-associated phenotypes detected by measures translatable to humans. *Proc Natl Acad Sci USA* 104(36):14501–14506.
- Hodgkinson CA, Goldman D, Jaeger J, Persaud S, Kane JM, Lipsky RH, Malhotra AK. 2004. Disrupted in schizophrenia 1 (DISC1): Association with schizophrenia, schizoaffective disorder, and bipolar disorder. *Am J Hum Genet* 75(5):862–872.
- Iwamoto K, Kakiuchi C, Bundo M, Ikeda K, Kato T. 2004. Molecular characterization of bipolar disorder by comparing gene expression profiles of postmortem brains of major mental disorders. *Mol Psychiatry* 9(4):406–416.
- Johansson C, Willeit M, Smedh C, Ekholm J, Paunio T, Kieseppa T, Lichtermann D, Praschak-Rieder N, Neumeister A, Nilsson LG, et al. 2003. Circadian clock-related polymorphisms in seasonal affective disorder and their relevance to diurnal preference. *Neuropsychopharmacology* 28(4):734–739.
- Johnson C, Drgon T, McMahon FJ, Uhl GR. 2009. Convergent genome wide association results for bipolar disorder and substance dependence. *Am J Med Genet Part B* 150B(2):182–190.
- Jurata LW, Bukhman YV, Charles V, Capriglione F, Bullard J, Lemire AL, Mohammed A, Pham Q, Laeng P, Brockman JA, et al. 2004. Comparison of microarray-based mRNA profiling technologies for identification of psychiatric disease and drug signatures. *J Neurosci Methods* 138(1–2):173–188.
- Kalman J, Palotas A, Juhasz A, Rimanoczy A, Hugyecz M, Kovacs Z, Galsi G, Szabo Z, Pakaski M, Feher LZ, et al. 2005. Impact of venlafaxine on gene expression profile in lymphocytes of the elderly with major depression—Evolution of antidepressants and the role of the “neuro-immune” system. *Neurochem Res* 30(11):1429–1438.
- Karege F, Schwald M, El Kouaissi R. 2004. Drug-induced decrease of protein kinase A activity reveals alteration in BDNF expression of bipolar affective disorder. *Neuropsychopharmacology* 29(4):805–812.
- Kelsoe JR, Niculescu AB III. 2002. Finding genes for bipolar disorder in the functional genomics era: From convergent functional genomics to phenomics and back. *CNS Spectr* 7(3):215–226.
- Kelsoe JR, Spence MA, Loetscher E, Foguet M, Sadovnick AD, Remick RA, Flodman P, Khristich J, Mroczkowski-Parker Z, Brown JL, et al. 2001. A genome survey indicates a possible susceptibility locus for bipolar disorder on chromosome 22. *Proc Natl Acad Sci USA* 98(2):585–590.
- Kerner B, Jasinska AJ, Deyoung J, Almonte M, Choi OW, Freimer NB. 2009. Polymorphisms in the GRIA1 gene region in psychotic bipolar disorder. *Am J Med Genet B Neuropsychiatr Genet*. 150B(1):24–32.
- Kim S, Choi KH, Baykiz AF, Gershenfeld HK. 2007. Suicide candidate genes associated with bipolar disorder and schizophrenia: An exploratory gene expression profiling analysis of post-mortem prefrontal cortex. *BMC Genomics* 8(1):413.
- Klempan TA, Sequeira A, Canetti L, Lalovic A, Ernst C, Ffrench-Mullen J, Turecki G. 2009. Altered expression of genes involved in ATP biosynthesis and GABAergic neurotransmission in the ventral prefrontal cortex of suicides with and without major depression. *Mol Psychiatry*. 14(2):175–189.
- Knable MB, Barci BM, Webster MJ, Meador-Woodruff J, Torrey EF. 2004. Molecular abnormalities of the hippocampus in severe psychiatric illness: Postmortem findings from the Stanley Neuropathology Consortium. *Mol Psychiatry* 9(6):609–620 544.
- Kurian SM, Le-Niculescu H, Patel SD, Bertram D, Davis J, Dike C, Yehyaw N, Lysaker P, Dustin J, Caligiuri M, et al. 2009. Identification of blood biomarkers for psychosis using convergent functional genomics. *Mol Psychiatry*. [Epub ahead of print]
- Lachman HM, Stopkova P, Papolos DF, Pedrosa E, Margolis B, Aghalar MR, Saito T. 2006. Analysis of synapsin III-196 promoter mutation in schizophrenia and bipolar disorder. *Neuropsychobiology* 53(2):57–62.
- Lachman HM, Pedrosa E, Petruolo OA, Cockerham M, Papolos A, Novak T, Papolos DF, Stopkova P. 2007. Increase in GSK3beta gene copy number variation in bipolar disorder. *Am J Med Genet Part B* 144B(3):259–265.
- Lahiri DK, Maloney B, Zawia NH. 2009. The LEARN model: An epigenetic explanation for idiopathic neurobiological diseases. *Mol Psychiatry* 14(11):992–1003.
- Lambert D, Middle F, Hamshere ML, Segurado R, Raybould R, Corvin A, Green E, O'Mahony E, Nikolov I, Mulcahy T, et al. 2005. Stage 2 of the Wellcome Trust UK-Irish bipolar affective disorder sibling-pair genome screen: Evidence for linkage on chromosomes 6q16-q21, 4q12-q21, 9p21, 10p14-p12 and 18q22. *Mol Psychiatry* 10(9):831–841.
- Le-Niculescu H, Balaraman Y, Patel S, Tan J, Sidhu K, Jerome RE, Edenberg HJ, Kuczenski R, Geyer MA, Nurnberger JI Jr, et al. 2007a. Towards understanding the schizophrenia code: An expanded convergent functional genomics approach. *Am J Med Genet Part B* 144B(2):129–158.
- Le-Niculescu H, McFarland MJ, Mamidipalli S, Ogden CA, Kuczenski R, Kurian SM, Salomon DR, Tsuang MT, Nurnberger JI Jr, Niculescu AB. 2007b. Convergent functional genomics of bipolar disorder: From animal model pharmacogenomics to human genetics and biomarkers. *Neurosci Biobehav Rev* 31(6):897–903.

- Le-Niculescu H, McFarland MJ, Ogden CA, Balaraman Y, Patel S, Tan J, Rodd ZA, Paulus M, Geyer MA, Edenberg HJ, et al. 2008. Phenomic, convergent functional genomic, and biomarker studies in a stress-reactive genetic animal model of bipolar disorder and co-morbid alcoholism. *Am J Med Genet Part B* 147B(2):134–166.
- Le-Niculescu H, Kurian SM, Yehyaw N, Dike C, Patel SD, Edenberg HJ, Tsuang MT, Salomon DR, Nurnberger JI Jr, Niculescu AB. 2009a. Identifying blood biomarkers for mood disorders using convergent functional genomics. *Mol Psychiatry* 14(2):156–174.
- Le-Niculescu H, Patel SD, Bhat M, Kuczenski R, Faraone SV, Tsuang MT, McMahon FJ, Schork NJ, Nurnberger JI Jr, Niculescu AB III. 2009b. Convergent functional genomics of genome-wide association data for bipolar disorder: Comprehensive identification of candidate genes, pathways and mechanisms. *Am J Med Genet Part B* 150B(2):155–181.
- Lenox RH, Gould TD, Manji HK. 2002. Endophenotypes in bipolar disorder. *Am J Med Genet* 114(4):391–406.
- Levinson DF, Evgrafov OV, Knowles JA, Potash JB, Weissman MM, Scheftner WA, Depaulo JR Jr, Crowe RR, Murphy-Eberenz K, Marta DH, et al. 2007. Genetics of recurrent early-onset major depression (GenRED): Significant linkage on chromosome 15q25-q26 after fine mapping with single nucleotide polymorphism markers. *Am J Psychiatry* 164(2):259–264.
- Lewohl JM, Wixey J, Harper CG, Dodd PR. 2005. Expression of MBP, PLP, MAG, CNP, and GFAP in the human alcoholic brain. *Alcohol Clin Exp Res* 29(9):1698–1705.
- Licinio J, Dong C, Wong ML. 2009. Novel sequence variations in the brain-derived neurotrophic factor gene and association with major depression and antidepressant treatment response. *Arch Gen Psychiatry* 66(5):488–497.
- Lin YM, Yang HC, Lai TJ, Fann CS, Sun HS. 2003. Receptor mediated effect of serotonergic transmission in patients with bipolar affective disorder. *J Med Genet* 40(10):781–786.
- Liu L, Foroud T, Xuei X, Berrettini W, Byerley W, Coryell W, El-Mallakh R, Gershon ES, Kelsoe JR, Lawson WB, et al. 2008. Evidence of association between brain-derived neurotrophic factor gene and bipolar disorder. *Psychiatr Genet* 18(6):267–274.
- Macgregor S, Visscher PM, Knott SA, Thomson P, Porteous DJ, Millar JK, Devon RS, Blackwood D, Muir WJ. 2004. A genome scan and follow-up study identify a bipolar disorder susceptibility locus on chromosome 1q42. *Mol Psychiatry* 9(12):1083–1090.
- Maeda K, Nwulia E, Chang J, Balkissoon R, Ishizuka K, Chen H, Zandi P, McClinnis MG, Sawa A. 2006. Differential expression of disrupted-in-schizophrenia (DISC1) in bipolar disorder. *Biol Psychiatry* 60(9):929–935.
- Magnusson A, Partonen T. 2005. The diagnosis, symptomatology, and epidemiology of seasonal affective disorder. *CNS Spectr* 10(8):625–634, quiz 1–14.
- Mansour HA, Wood J, Logue T, Chowdari KV, Dayal M, Kupfer DJ, Monk TH, Devlin B, Nimgaonkar VL. 2006. Association study of eight circadian genes with bipolar I disorder, schizoaffective disorder and schizophrenia. *Genes Brain Behav* 5(2):150–157.
- Matigian N, Windus L, Smith H, Filippich C, Pantelis C, McGrath J, Mowry B, Hayward N. 2007. Expression profiling in monozygotic twins discordant for bipolar disorder reveals dysregulation of the WNT signalling pathway. *Mol Psychiatry* 12(9):815–825.
- Maziade M, Roy MA, Chagnon YC, Cliche D, Fournier JP, Montgrain N, Dion C, Lavallee JC, Garneau Y, Gingras N, et al. 2005. Shared and specific susceptibility loci for schizophrenia and bipolar disorder: A dense genome scan in Eastern Quebec families. *Mol Psychiatry* 10(5):486–499.
- McAuley EZ, Fullerton JM, Blair IP, Donald JA, Mitchell PB, Schofield PR. 2009. Association between the serotonin 2A receptor gene and bipolar affective disorder in an Australian cohort. *Psychiatr Genet* 9(5):244–252.
- McClung CA. 2007. Circadian genes, rhythms and the biology of mood disorders. *Pharmacol Ther* 114(2):222–232.
- McGrath CL, Glatt SJ, Sklar P, Le-Niculescu H, Kuczenski R, Doyle AE, Biederman J, Mick E, Faraone SV, Niculescu AB, et al. 2009. Evidence for genetic association of RORB with bipolar disorder. *BMC Psychiatry* 9:70.
- McInnes LA, Escamilla MA, Service SK, Reus VI, Leon P, Silva S, Rojas E, Spesny M, Baharloo S, Blankenship K, et al. 1996. A complete genome screen for genes predisposing to severe bipolar disorder in two Costa Rican pedigrees. *Proc Natl Acad Sci USA* 93(23):13060–13065.
- Middleton FA, Pato CN, Gentile KL, McGann L, Brown AM, Trauzzi M, Diab H, Morley CP, Medeiros H, Macedo A, et al. 2005. Gene expression analysis of peripheral blood leukocytes from discordant sib-pairs with schizophrenia and bipolar disorder reveals points of convergence between genetic and functional genomic approaches. *Am J Med Genet Part B* 136B(1):12–25.
- Millar JK, Mackie S, Clapcote SJ, Murdoch H, Pickard BS, Christie S, Muir WJ, Blackwood DH, Roder JC, Houslay MD, et al. 2007. Disrupted in schizophrenia 1 and phosphodiesterase 4B: Towards an understanding of psychiatric illness. *J Physiol* 584(Pt2):401–405.
- Morissette J, Villeneuve A, Bordeleau L, Rochette D, Laberge C, Gagne B, Laprise C, Bouchard G, Plante M, Gobeil L, et al. 1999. Genome-wide search for linkage of bipolar affective disorders in a very large pedigree derived from a homogeneous population in quebec points to a locus of major effect on chromosome 12q23-q24. *Am J Med Genet* 88(5):567–587.
- Nakatani N, Hattori E, Ohnishi T, Dean B, Iwayama Y, Matsumoto I, Kato T, Osumi N, Higuchi T, Niwa S, et al. 2006. Genome-wide expression analysis detects eight genes with robust alterations specific to bipolar I disorder: Relevance to neuronal network perturbation. *Hum Mol Genet* 15(12):1949–1962.
- Neves-Pereira M, Mundo E, Muglia P, King N, Macciardi F, Kennedy JL. 2002. The brain-derived neurotrophic factor gene confers susceptibility to bipolar disorder: Evidence from a family-based association study. *Am J Hum Genet* 71(3):651–655.
- Niculescu AB III. 2006. Polypharmacy in oligopopulations: What psychiatric genetics can teach biological psychiatry. *Psychiatr Genet* 16(6):241–244.
- Niculescu AB III, Kelsoe JR. 2001. Convergent functional genomics: Application to bipolar disorder. *Ann Med* 33(4):263–271.
- Niculescu AB, Le-Niculescu H. 2010. Convergent functional genomics: What we have learned and can learn about genes, pathways, and mechanisms. *Neuropsychopharmacology* 35(1):355–356.
- Niculescu A, Segal D, Kuczenski R, Barrett T, Hauger R, Kelsoe J. 2000a. Identifying a series of candidate genes for mania and psychosis: A convergent functional genomics approach. *Physiol Genomics* 4(1):83–91.
- Niculescu AB III, Segal DS, Kuczenski R, Barrett T, Hauger RL, Kelsoe JR. 2000b. Identifying a series of candidate genes for mania and psychosis: A convergent functional genomics approach. *Physiol Genomics* 4(1):83–91.
- Niculescu AB, Lulow LL, Ogden CA, Le-Niculescu H, Salomon DR, Schork NJ, Caligiuri MP, Lohr JB. 2006. PhenoChipping of psychotic disorders: A novel approach for deconstructing and quantitating psychiatric phenotypes. *Am J Med Genet Part B* 141B(6):653–662.
- Niculescu AB III, Schork NJ, Salomon DR. 2009. Mindscape: A convergent perspective on life, mind, consciousness and happiness. *J Affect Disord*

- Nievergelt CM, Kripke DF, Barrett TB, Burg E, Remick RA, Sadvnick AD, McElroy SL, Keck PE Jr, Schork NJ, Kelsoe JR. 2006. Suggestive evidence for association of the circadian genes PERIOD3 and ARNTL with bipolar disorder. *Am J Med Genet Part B* 141B(3):234–241.
- Novak G, Seeman P, Tallerico T. 2006. Increased expression of calcium/calmodulin-dependent protein kinase IIbeta in frontal cortex in schizophrenia and depression. *Synapse* 59(1):61–68.
- Numata S, Iga J, Nakataki M, Tayoshi S, Taniguchi K, Sumitani S, Tomotake M, Tanahashi T, Itakura M, Kamegaya Y, et al. 2009. Gene expression and association analyses of the phosphodiesterase 4B (PDE4B) gene in major depressive disorder in the Japanese population. *Am J Med Genet Part B* 150B(4):527–534.
- Nunes PV, Forlenza OV, Gattaz WF. 2007. Lithium and risk for Alzheimer's disease in elderly patients with bipolar disorder. *Br J Psychiatry* 190:359–360.
- O'Donovan MC, Craddock NJ, Owen MJ. 2009. Genetics of psychosis: insights from views across the genome. *Hum Genet* 126(1):3–12.
- Ogden CA, Rich ME, Schork NJ, Paulus MP, Geyer MA, Lohr JB, Kuczenski R, Niculescu AB. 2004. Candidate genes, pathways and mechanisms for bipolar (manic-depressive) and related disorders: An expanded convergent functional genomics approach. *Mol Psychiatry* 9(11):1007–1029.
- Oh ES, Savonenko AV, King JF, Fangmark Tucker SM, Rudow GL, Xu G, Borchelt DR, Troncoso JC. 2008. Amyloid precursor protein increases cortical neuron size in transgenic mice. *Neurobiol Aging*. 30(8):1238–1244.
- Padmos RC, Hillegers MH, Knijff EM, Vonk R, Bouvy A, Staal FJ, de Ridder D, Kupka RW, Nolen WA, Drexhage HA. 2008. A discriminating messenger RNA signature for bipolar disorder formed by an aberrant expression of inflammatory genes in monocytes. *Arch Gen Psychiatry* 65(4):395–407.
- Park N, Joo SH, Cheng R, Liu J, Loth JE, Lilliston B, Nee J, Grunn A, Kanyas K, Lerer B, et al. 2004. Linkage analysis of psychosis in bipolar pedigrees suggests novel putative loci for bipolar disorder and shared susceptibility with schizophrenia. *Mol Psychiatry* 9(12):1091–1099.
- Parlapani E, Schmitt A, Erdmann A, Bernstein HG, Breunig B, Gruber O, Petroianu G, von Wilmsdorff M, Schneider-Axmann T, Honer W, et al. 2009. Association between myelin basic protein expression and left entorhinal cortex pre-alpha cell layer disorganization in schizophrenia. *Brain Res*.
- Partonen T, Treutlein J, Alpman A, Frank J, Johansson C, Depner M, Aron L, Rietschel M, Wellek S, Soronen P, et al. 2007. Three circadian clock genes *Per2*, *Arntl*, and *Npas2* contribute to winter depression. *Ann Med* 39(3):229–238.
- Paynter NP, Chasman DI, Pare G, Buring JE, Cook NR, Miletich JP, Ridker PM. 2010. Association between a literature-based genetic risk score and cardiovascular events in women. *JAMA* 303(7):631–637.
- Pennington K, Beasley CL, Dicker P, Fagan A, English J, Pariante CM, Wait R, Dunn MJ, Cotter DR. 2008. Prominent synaptic and metabolic abnormalities revealed by proteomic analysis of the dorsolateral prefrontal cortex in schizophrenia and bipolar disorder. *Mol Psychiatry*. 13(12):1102–1117.
- Perlis RH, Purcell S, Fagerness J, Kirby A, Petryshen TL, Fan J, Sklar P. 2008. Family-based association study of lithium-related and other candidate genes in bipolar disorder. *Arch Gen Psychiatry* 65(1):53–61.
- Pezawas L, Meyer-Lindenberg A, Goldman AL, Verchinski BA, Chen G, Kolachana BS, Egan MF, Mattay VS, Hariri AR, Weinberger DR. 2008. Evidence of biologic epistasis between BDNF and SLC6A4 and implications for depression. *Mol Psychiatry* 13(7):654, 709–716.
- Pillai A. 2008. Decreased expression of Sprouty2 in the dorsolateral prefrontal cortex in schizophrenia and bipolar disorder: A correlation with BDNF expression. *PLoS ONE* 3(3):e1784.
- Pletnikov MV, Xu Y, Ovanesov MV, Kamiya A, Sawa A, Ross CA. 2007. PC12 cell model of inducible expression of mutant DISC1: New evidence for a dominant-negative mechanism of abnormal neuronal differentiation. *Neurosci Res* 58(3):234–244.
- Potash JB, Zandi PP, Willour VL, Lan TH, Huo Y, Avramopoulos D, Shugart YY, MacKinnon DF, Simpson SG, McMahon FJ, et al. 2003. Suggestive linkage to chromosomal regions 13q31 and 22q12 in families with psychotic bipolar disorder. *Am J Psychiatry* 160(4):680–686.
- Potash JB, Buervenich S, Cox NJ, Zandi PP, Akula N, Steele J, Rathe JA, Avramopoulos D, Detera-Wadleigh SD, Gershon ES, et al. 2008. Gene-based SNP mapping of a psychotic bipolar affective disorder linkage region on 22q12.3: Association with HMG2L1 and TOM1. *Am J Med Genet Part B* 147B(1):59–67.
- Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, Sklar P. 2009. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 460(7256):748–752.
- Radhakrishna U, Senol S, Herken H, Gucuyener K, Gehrig C, Blouin JL, Akarsu NA, Antonarakis SE. 2001. An apparently dominant bipolar affective disorder (BPAD) locus on chromosome 20p11.2–q11.2 in a large Turkish pedigree. *Eur J Hum Genet* 9(1):39–44.
- Ranade SS, Mansour H, Wood J, Chowdari KV, Brar LK, Kupfer DJ, Nimgaonkar VL. 2003. Linkage and association between serotonin 2A receptor gene polymorphisms and bipolar I disorder. *Am J Med Genet Part B* 121B(1):28–34.
- Rice TK, Schork NJ, Rao DC. 2008. Methods for handling multiple testing. *Adv Genet* 60:293–308.
- Ripperger JA, Schibler U. 2006. Rhythmic CLOCK-BMAL1 binding to multiple E-box motifs drives circadian Dbp transcription and chromatin transitions. *Nat Genet* 38(3):369–374.
- Rodd ZA, Bertsch BA, Strother WN, Le-Niculescu H, Balaraman Y, Hayden E, Jerome RE, Lumeng L, Nurnberger Jr, Edenberg HJ, et al. 2007. Candidate genes, pathways and mechanisms for alcoholism: An expanded convergent functional genomics approach. *Pharmacogenomics J* 7(4):222–256.
- Rohn TT, Vyas V, Hernandez-Estrada T, Nichol KE, Christie LA, Head E. 2008. Lack of pathology in a triple transgenic mouse model of Alzheimer's disease after overexpression of the anti-apoptotic protein Bcl-2. *J Neurosci* 28(12):3051–3059.
- Ryan MM, Lockstone HE, Huffaker SJ, Wayland MT, Webster MJ, Bahn S. 2006. Gene expression analysis of bipolar disorder reveals downregulation of the ubiquitin cycle and alterations in synaptic genes. *Mol Psychiatry* 11(10):965–978.
- Sawamura N, Sawamura-Yamamoto T, Ozeki Y, Ross CA, Sawa A. 2005. A form of DISC1 enriched in nucleus: Altered subcellular distribution in orbitofrontal cortex in psychosis and substance/alcohol abuse. *Proc Natl Acad Sci USA* 102(4):1187–1192.
- Schosser A, Gaysina D, Cohen-Woods S, Chow PC, Martucci L, Craddock N, Farmer A, Korszun A, Gunasinghe C, Gray J, et al. 2009. Association of DISC1 and TSNAX genes and affective disorders in the depression case-control (DeCC) and bipolar affective case-control (BACCS) studies. *Mol Psychiatry*. [Epub ahead of print]
- Schulze TG, Chen YS, Badner JA, McInnis MG, DePaulo JR Jr, McMahon FJ. 2003. Additional, physically ordered markers increase linkage signal for bipolar disorder on chromosome 18q22. *Biol Psychiatry* 53(3):239–243.
- Schulze TG, Detera-Wadleigh SD, Akula N, Gupta A, Kassem L, Steele J, Pearl J, Strohmaier J, Breuer R, Schwarz M, et al. 2009. Two variants in

- Ankyrin 3 (ANK3) are independent genetic risk factors for bipolar disorder. *Mol Psychiatry* 14(5):487–491.
- Segurado R, Detera-Wadleigh SD, Levinson DF, Lewis CM, Gill M, Nurnberger JI Jr, Craddock N, DePaulo JR, Baron M, Gershon ES, et al. 2003. Genome scan meta-analysis of schizophrenia and bipolar disorder, part III: Bipolar disorder. *Am J Hum Genet* 73(1):49–62.
- Sen S, Duman R, Sanacora G. 2008. Serum brain-derived neurotrophic factor, depression, and antidepressant medications: Meta-analyses and implications. *Biol Psychiatry*
- Sequeira A, Klempan T, Canetti L, Ffrench-Mullen J, Benkelfat C, Rouleau GA, Turecki G. 2007. Patterns of gene expression in the limbic system of suicides with and without major depression. *Mol Psychiatry* 12(7): 640–655.
- Shi J, Wittke-Thompson JK, Badner JA, Hattori E, Potash JB, Willour VL, McMahon FJ, Gershon ES, Liu C. 2008. Clock genes may influence bipolar disorder susceptibility and dysfunctional circadian rhythm. *Am J Med Genet B Neuropsychiatr Genet*. 147B(7):1047–1055.
- Sklar P, Gabriel SB, McInnis MG, Bennett P, Lim YM, Tsan G, Schaffner S, Kirov G, Jones I, Owen M, et al. 2002. Family-based association study of 76 candidate genes in bipolar disorder: BDNF is a potential risk locus. Brain-derived neurotrophic factor. *Mol Psychiatry* 7(6):579–593.
- Sklar P, Pato MT, Kirby A, Petryshen TL, Medeiros H, Carvalho C, Macedo A, Dourado A, Coelho I, Valente J, et al. 2004. Genome-wide scan in Portuguese Island families identifies 5q31-5q35 as a susceptibility locus for schizophrenia and psychosis. *Mol Psychiatry* 9(2):213–218.
- Sklar P, Smoller JW, Fan J, Ferreira MAR, Perlis RH, Chambert K, Nimgaonkar VL, McQueen MB, Faraone SV, Kirby A, et al. 2008. Whole-genome association study of bipolar disorder. *Mol Psychiatry* 13(6):558–569.
- Smith EN, Bloss CS, Badner JA, Barrett T, Belmonte PL, Berrettini W, Byerley W, Coryell W, Craig D, Edenberg HJ, et al. 2009. Genome-wide association study of bipolar disorder in European American and African American individuals. *Mol Psychiatry* 14(8):755–763.
- Sokolov BP, Polesskaya OO, Uhl GR. 2003. Mouse brain gene expression changes after acute and chronic amphetamine. *J Neurochem* 84(2): 244–252.
- Stumpf MP, Thorne T, de Silva E, Stewart R, An HJ, Lappe M, Wiuf C. 2008. Estimating the size of the human interactome. *Proc Natl Acad Sci USA* 105(19):6959–6964.
- Sun X, Steffens DC, Au R, Folstein M, Summergrad P, Yee J, Rosenberg I, Mwamburi DM, Qiu WQ. 2008. Amyloid-associated depression: A prodromal depression of Alzheimer disease? *Arch Gen Psychiatry* 65(5):542–550.
- Szczepankiewicz A, Rybakowski JK, Suwalska A, Skibinska M, Leszczynska-Rodziewicz A, Dmitrzak-Weglarz M, Czerski PM, Hauser J. 2006. Association study of the glycogen synthase kinase-3beta gene polymorphism with prophylactic lithium response in bipolar patients. *World J Biol Psychiatry* 7(3):158–161.
- Takahashi JS, Shimomura K, Kumar V. 2008. Searching for genes underlying behavior: Lessons from circadian rhythms. *Science* 322(5903): 909–912.
- Tan HY, Callicott JH, Weinberger DR. 2008. Intermediate phenotypes in schizophrenia genetics redux: Is it a no brainer? *Mol Psychiatry* 13(3):233–238.
- Thomson PA, Wray NR, Millar JK, Evans KL, Hellard SL, Condie A, Muir WJ, Blackwood DH, Porteous DJ. 2005. Association between the TRAX/DISC locus and both bipolar disorder and schizophrenia in the Scottish population. *Mol Psychiatry* 10(7):657–668, 616.
- Thomson PA, Christoforou A, Morris SW, Adie E, Pickard BS, Porteous DJ, Muir WJ, Blackwood DH, Evans KL. 2007. Association of Neuregulin 1 with schizophrenia and bipolar disorder in a second cohort from the Scottish population. *Mol Psychiatry* 12(1):94–104.
- Tkachev D, Mimmack ML, Ryan MM, Wayland M, Freeman T, Jones PB, Starkey M, Webster MJ, Yolken RH, Bahn S. 2003. Oligodendrocyte dysfunction in schizophrenia and bipolar disorder. *Lancet* 362(9386): 798–805.
- Tochigi M, Iwamoto K, Bundo M, Sasaki T, Kato N, Kato T. 2008. Gene expression profiling of major depression and suicide in the prefrontal cortex of postmortem brains. *Neurosci Res* 60(2):184–191.
- Torrey EF, Barci BM, Webster MJ, Bartko JJ, Meador-Woodruff JH, Knable MB. 2005. Neurochemical markers for schizophrenia, bipolar disorder, and major depression in postmortem brains. *Biol Psychiatry* 57(3): 252–260.
- Uher R, Huezio-Diaz P, Perroud N, Smith R, Rietschel M, Mors O, Hauser J, Maier W, Kozel D, Henigsberg N, et al. 2009. Genetic predictors of response to antidepressants in the GENDEP project. *Pharmacogenomics J*
- van der Veen DR, Minh NL, Gos P, Arneric M, Gerkema MP, Schibler U. 2006. Impact of behavior on central and peripheral circadian clocks in the common vole *Microtus arvalis*, a mammal with ultradian rhythms. *Proc Natl Acad Sci USA* 103(9):3393–3398.
- van West D, Van Den Eede F, Del-Favero J, Souery D, Norrback KF, Van Duijn C, Sluijs S, Adolfsson R, Mendlewicz J, Deboutte D, et al. 2006. Glucocorticoid receptor gene-based SNP analysis in patients with recurrent major depression. *Neuropsychopharmacology* 31(3):620–627.
- Vawter MP, Thatcher L, Usen N, Hyde TM, Kleinman JE, Freed WJ. 2002. Reduction of synapsin in the hippocampus of patients with bipolar disorder and schizophrenia. *Mol Psychiatry* 7(6):571–578.
- Vawter MP, Tomita H, Meng F, Bolstad B, Li J, Evans S, Choudary P, Atz M, Shao L, Neal C, et al. 2006. Mitochondrial-related gene expression changes are sensitive to agonal-pH state: Implications for brain disorders. *Mol Psychiatry*
- Wager-Smith K, Kay SA. 2000. Circadian rhythm genetics: From flies to mice to humans. *Nat Genet* 26(1):23–27.
- Wakabayashi Y, Uchida S, Funato H, Matsubara T, Watanuki T, Otsuki K, Fujimoto M, Nishida A, Watanabe Y. 2008. State-dependent changes in the expression levels of NCAM-140 and L1 in the peripheral blood cells of bipolar disorders, but not in the major depressive disorders. *Prog Neuropsychopharmacol Biol Psychiatry* 32(5):1199–1205.
- Walss-Bass C, Raventos H, Montero AP, Armas R, Dassori A, Contreras S, Liu W, Medina R, Levinson DF, Pereira M, et al. 2006. Association analyses of the neuregulin 1 gene with schizophrenia and manic psychosis in a Hispanic population. *Acta Psychiatr Scand* 113(4):314–321.
- Wirz-Justice A. 2006. Biological rhythm disturbances in mood disorders. *Int Clin Psychopharmacol* 21(Suppl. 1): S11–S15.
- Wirz-Justice A, Terman M, Oren DA, Goodwin FK, Kripke DF, Whybrow PC, Wisner KL, Wu JC, Lam RW, Berger M, et al. 2004. Brightening depression. *Science* 303(5657):467–469.
- Wong ML, Whelan F, Deloukas P, Whittaker P, Delgado M, Cantor RM, McCann SM, Licinio J. 2006. Phosphodiesterase genes are associated with susceptibility to major depression and antidepressant treatment response. *Proc Natl Acad Sci USA* 103(41):15124–15129.
- Wong ML, Dong C, Maestre-Mesa J, Licinio J. 2008. Polymorphisms in inflammation-related genes are associated with susceptibility to major depression and antidepressant response. *Mol Psychiatry* 13(8):800–812.
- WTCC. 2007. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* 447(7145):661–678.
- Xing G, Russell S, Hough C, O'Grady J, Zhang L, Yang S, Zhang LX, Post R. 2002. Decreased prefrontal CaMKII alpha mRNA in bipolar illness. *Neuroreport* 13(4):501–505.

- Yang S, Van Dongen HP, Wang K, Berrettini W, Bucan M. 2008. Assessment of circadian function in fibroblasts of patients with bipolar disorder. *Mol Psychiatry*
- Zamvil S, Nelson P, Trotter J, Mitchell D, Knobler R, Fritz R, Steinman L. 1985. T-cell clones specific for myelin basic protein induce chronic relapsing paralysis and demyelination. *Nature* 317(6035):355–358.
- Zhang EE, Liu AC, Hirota T, Miraglia LJ, Welch G, Pongsawakul PY, Liu X, Atwood A, Huss JW III, Janes J, et al. 2009. A genome-wide RNAi screen for modifiers of the circadian clock in human cells. *Cell* 139(1):199–210.
- Zubenko GS, Hughes HB, Stiffler JS, Zubenko WN, Kaplan BB. 2002. Genome survey for susceptibility loci for recurrent, early-onset major depression: Results at 10cM resolution. *Am J Med Genet* 114(4):413–422.
- Zubenko GS, Maher B, Hughes HB III, Zubenko WN, Stiffler JS, Kaplan BB, Marazita ML. 2003. Genome-wide linkage survey for genetic loci that influence the development of depressive disorders in families with recurrent, early-onset, major depression. *Am J Med Genet Part B* 123B(1):1–18.



Supplementary Information

**Figure 1S. CFG of GWAS for Bipolar Disorder
-cross-matching genetic with gene expression data**

3 GWAS (WTCC, NIMH, German)	1565 genes
+ Step-BD	+92 genes
+ GAIN-BP	+169 genes
Total	1826 genes

Up to 10% of the genome may be involved
directly or indirectly in mood regulation

Figure 2S: Gene size vs. CFG score.




Figure 3S: Two-Dimensional Genetic Risk Prediction Matrix

Genetic Risk Prediction Panel																	
	Subject	Gene 1								--	--	--	Gene 56				Genetic Risk Prediction Score (0-100) (sum of allele scores/ number of alleles) x 100
		SNP1		SNP2		SNP3		SNP4		--	--	--	SNP1		SNP2		
		Allele 1	Allele 2	Allele 1	Allele 2	Allele 1	Allele 2	Allele 1	Allele 2	--	--	--	Allele 1	Allele 2	Allele 1	Allele 2	
Bipolar	1	1	1	1	0	1	0	0	1	--	--	--	0	1	1	0	51.709
	2	1	1	0	1	1	1	1	1	--	--	--	1	0	1	1	57.265
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	406	0	1	1	1	0	0	1	0	--	--	--	0	0	0	0	55.128
407	1	0	0	1	1	1	1	1	--	--	--	1	0	1	1	55.508	
Control	1	0	1	0	0	1	1	0	1	--	--	--	1	1	1	1	50.000
	2	1	0	0	1	1	0	0	0	--	--	--	1	0	1	0	47.863
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	1033	0	0	1	0	0	1	0	1	--	--	--	0	1	0	1	49.576
1034	0	0	0	0	1	1	0	1	--	--	--	1	1	1	1	46.186	

Table 1S. Demographics of the randomly split cohorts from GAIN-BP. NA- not available.

	European American (EA)				African American (AA)			
	2/3 Split Sample		1/3 Split Sample		2/3 Split Sample		1/3 Split Sample	
	Bipolar	Control	Bipolar	Control	Bipolar	Control	Bipolar	Control
Number of Subjects (males: females)	272 (133: 139)	690 (356: 334)	135 (64: 71)	344 (178: 166)	212 (67: 145)	448 (182: 266)	105 (33: 72)	223 (90: 133)
Age mean years (SD) range	43.32 (13.62) 20 to 82	51.41 (17.67) 18 to 90	43.19 (13.05) 21 to 77	53.94 (17.27) 18 to 84	42.92 (10.41) 17 to 70	NA	39.91 (10.83) 19 to 61	NA
Episode Frequency mean (SD) range	1.50 (5.04) 0.06 to 42.60	--	1.49 (2.20) 0.06 to 6.97	--	2.01 (4.94) 0.02 to 50.05	--	1.72 (2.65) 0.02 to 14.71	--

L. Suppl. Note. Members of the Bipolar Disorder Genetics (BiGS) Consortium

E N Smith^{1,2,24}, C S Bloss^{1,3,24}, J A Badner⁴, T Barrett⁵, P L Belmonte⁶, W Berrettini⁷, W Byerley⁸, W Coryell⁹, D Craig¹⁰, H J Edenberg^{11,12}, E Eskin¹³, T Foroud¹², E S Gershon⁴, T A Greenwood¹⁴, M Hipolito¹⁵, D L Koller¹⁶, W B Lawson¹⁵, C Liu⁴, F Lohoff⁷, M G McInnis¹⁷, F J McMahon¹⁸, D B Mirel¹⁹, S S Murray^{1,2,3}, C Nievergelt¹⁴, J Nurnberger¹⁶, E A Nwulia¹⁵, J Paschall²⁰, J B Potash⁶, J Rice²¹, T G Schulze¹⁸, W Scheftner²², C Panganiban¹⁰, N Zaitlen¹³, P P Zandi⁶, S Zöllner¹⁷, N J Schork^{1,2,3}, J R Kelsøe^{14,23}

¹Scripps Genomic Medicine, Scripps Translational Science Institute, La Jolla, CA, USA;

²Department of Molecular and Experimental Medicine, The Scripps Research Institute, La Jolla, CA, USA;

³Scripps Health, La Jolla, CA, USA; ⁴Department of Psychiatry, University of Chicago, Chicago, IL, USA;

⁵Department of Psychiatry, Portland VA Medical Center, Portland, OR, USA;

⁶Department of Psychiatry, Johns Hopkins School of Medicine, Baltimore, MD, USA;

⁷Department of Psychiatry, University of Pennsylvania, Philadelphia, PA, USA; ⁸Department of Psychiatry, University of California, San Francisco, San Francisco, CA, USA;

⁹Department of Psychiatry, University of Iowa, Iowa City, IA, USA; ¹⁰Neurogenomics Division, The Translational Genomics Research Institute, Phoenix, AZ, USA;

¹¹Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN, USA;

¹²Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN, USA;

¹³Department of Computer Science, University of California, Los Angeles, Los Angeles, CA, USA;

¹⁴Department of Psychiatry, University of California, San Diego, La Jolla, CA, USA;

¹⁵Department of Psychiatry, Howard University, Washington, DC, USA;

¹⁶Department of Psychiatry, Indiana University School of Medicine, Indianapolis, IN, USA;

¹⁷Department of Psychiatry, University of Michigan, Ann Arbor, MI, USA; ¹⁸Genetic Basis of Mood and Anxiety Disorders Unit, National Institute of Mental Health Intramural Research Program, National Institutes of Health, US Department of Health and Human Services, Bethesda, MD, USA;

¹⁹Broad Institute of Harvard University and Massachusetts Institute of Technology, Cambridge, MA, USA;

²⁰National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD, USA;

²¹Division of Biostatistics, Washington University, St Louis, MO, USA;

²²Department of Psychiatry, Rush University, Chicago, IL, USA;

²³Department of Psychiatry, VA San Diego Healthcare System, La Jolla, CA, USA

USA